

Rec'd PCT/PTO 16 JUN 2005 539 868

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



10/539868



(43) International Publication Date
5 August 2004 (05.08.2004)

PCT

(10) International Publication Number
WO 2004/064601 A2

(51) International Patent Classification:

A61B

(74) Agents: KELLY, Beth, L. et al.; Two Embarcadero Center, 8th Floor, San Francisco, CA 94111 (US).

(11) International Application Number:

PCT/US2004/001153

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 15 January 2004 (15.01.2004)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10/346,367

15 January 2003 (15.01.2003)

US

(71) Applicant (for all designated States except US): THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 1111 Franklin Street, 12th Floor, Oakland, CA 94607 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SLAMON, Dennis, J. [US/US]; 23122 Calvert Street, Woodland Hills, CA 91367 (US). ANDERSON, Lee, A. [US/US]; 1134 Cardiff Avenue, Los Angeles, CA 90035 (US). GINTHER, Charles, L. [US/US]; 1134 Cardiff Avenue, Los Angeles, CA 90035 (US).


Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

[Continued on next page]

(54) Title: AMPLIFIED AND OVEREXPRESSED GENE IN COLORECTAL CANCERS

RNA and protein expression of 26#77 is higher in paired colon tumors vs. normals



(57) Abstract: This invention relates to methods to diagnose colon cancer and other proliferative diseases by detecting amplification of genes and gene products on chromosome 20q13.2, a region whose amplification is associated with a poor cancer prognosis.

BEST AVAILABLE COPY

WO 2004/064601 A2



Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

AMPLIFIED AND OVEREXPRESSED GENE IN COLORECTAL CANCERS

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application is a continuation in part of and claims the benefit of priority of U.S.
5 Patent Application No. 10/346,367, filed on January 15, 2003, which is herein incorporated
by reference for all purposes.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

10 [0002] This invention was made with Government support by Grant No. CA32737,
awarded by the National Institutes of Health. The Government has certain rights in this
invention.

FIELD OF THE INVENTION

[0003] This invention relates to methods to diagnose colon cancer and other proliferative
15 diseases.

BACKGROUND OF THE INVENTION

[0004] Chromosome abnormalities are often associated with genetic disorders,
degenerative diseases, and cancer. The deletion or multiplication of copies of whole
chromosomes and the deletion or amplifications of chromosomal segments or specific
20 regions are common occurrences in cancer (Smith (1991) *Breast Cancer Res. Treat.* 18:
Suppl. 1:5-14; van de Vijer (1991) *Biochim. Biophys. Acta.* 1072:33-50). In fact,
amplifications and deletions of DNA sequences can be the cause of a cancer. For example,
proto-oncogenes and tumor-suppressor genes, respectively, are frequently characteristic of
tumorigenesis (Dutrillaux (1990) *Cancer Genet. Cytogenet.* 49: 203-217). Clearly, the
25 identification and cloning of specific genomic regions associated with cancer is crucial both
to the study of tumorigenesis and in developing better means of diagnosis and prognosis.

[0005] One of the amplified regions found in studies of breast and colon cancer cells is on
chromosome 20, specifically, 20q13.2 (*see, e.g.* WO98/02539). Amplification of 20q13.2

was subsequently found to occur in a variety of tumor types and to be associated with aggressive tumor behavior. Increased 20q13.2 copy number has been found in 40% of breast cancer cell lines and 18% of primary breast tumors (Kalliioniemi (1994) *Proc. Natl. Acad. Sci. USA* 91: 2156-2160). Copy number gains at 20q13.2 have also been reported in greater than 25% of cancers of the ovary (Iwabuchi (1995) *Cancer Res.* 55:6172-6180), colon (Schlegel (1995) *Cancer Res.* 55: 6002-6005 and WO02/06526), head-and-neck (Bockmuhl (1996) *Laryngor.* 75: 408-414), brain (Mohapatra (1995) *Genes Chromosomes Cancer* 13: 86-93), pancreas (Solinas-Toldo (1996) *Genes Chromosomes Cancer* 20:399-407).

[0006] A number of studies have elucidated genetic alterations that occur during the development of colorectal tumors. For instance, deletions of p53 genes on chromosome 17p are often late events associated with the transition from the benign (adenoma) to the malignant (carcinoma) state. See Vogelstein *et al.*, *New England Journal of Medicine*, 319:525 (1988), Fearon and Vogelstein, *Cell*, 61:759-767 (1990) and Baker *et al.* *Cancer Res.* 50:7717-22 (1990). More recently, comparative genomic hybridization has shown that specific patterns of chromosomal gains and losses take place during colorectal carcinogenesis (see, e.g. Schlegel, *et al.* *Cancer Research.* 55, 6002-6005 (1995); Ried, *et al.* *Genes, Chromosomes & Cancer* 15, 234-245 (1996); and Nakao *et al.*, *Jpn. J. Surg.* 28, 567-569 (1998). These changes included overrepresentation (amplification) of large portion of chromosome 20 material.

[0007] The identification of new genes that are responsible for carcinogenesis is obviously great use in diagnosis, prognosis and treatment of these diseases. The present invention addresses these and other needs.

BRIEF SUMMARY OF THE INVENTION

[0008] This invention provides a method for determining the presence or absence of a colorectal cancer cell in a patient, by determining the level of a target nucleic acid that encodes the 26#77 protein (e.g., SEQ ID NO: 2) in a biological sample from the patient. In one embodiment, the target nucleic acid comprises a sequence at least 80% identical to SEQ ID NO: 1. In a further embodiment, the biological sample can include isolated nucleic acids. In another embodiment, the nucleic acids are amplified before the level of the target nucleic acid is determined. In an additional embodiment the isolated nucleic acids are mRNA.

[0009] The invention also provides a method for determining the presence or absence of a colorectal cancer cell in a patient, by determining the level of a target nucleic acid that encodes the Copine 1 (CPNE 1) protein, the Integrin B4 binding protein (ITGB4BP), RNA Export homolog (RAE1), bone morphogenic protein 7 (BMP7), G protein, alpha stimulating activity polypeptide 1 (GNAS), eukaryotic translation initiation factor 2, subunit 2 beta (EIF2S2), dynein light chain A2 (DNCL2A), proteasome subunit α -7 (PSMA7), activity dependent neuroprotector (ADNP), C20orf129, C20orf52, C20orf20, or C20orf188 (*e.g.*, SEQ ID NO:4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, or 28) in a biological sample from the patient. In one embodiment, the target nucleic acid comprises a sequence at least 80% identical to SEQ ID NO:3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, or 27. In a further embodiment, the biological sample can include isolated nucleic acids. In another embodiment, the nucleic acids are amplified before the level of the target nucleic acid is determined. In an additional embodiment the isolated nucleic acids are mRNA.

[0010] In one aspect, the biological sample is colorectal tissue and the step of determining the level of target nucleic acid is carried out using in situ hybridization.

[0011] In another aspect, the step of determining the level of target nucleic acid is carried out using a labeled nucleic acid probe that selectively hybridizes to SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, or 27 under stringent hybridization conditions. The nucleic acid probe can be immobilized to a solid support. In a further aspect, the step of determining the level of target nucleic acid is carried out using Northern blot analysis.

[0012] In one embodiment, the step of determining the level of the target nucleic acid is carried out by comparing the amount of the target nucleic acid in the biological sample to the amount of the target nucleic acid in a reference sample. The reference sample can be from normal colorectal tissue.

[0013] In another embodiment, the levels of 26#77 encoding nucleic acid are determined when the patient is undergoing a therapeutic regimen to treat colorectal cancer. The levels of 26#77 encoding nucleic acid can also be determined when the patient is suspected of having colorectal cancer. Similarly, levels of CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 encoding nucleic acid are determined when the patient is undergoing a therapeutic regimen to treat colorectal cancer, and can also be determined when the patient is suspected of having colorectal cancer.

[0014] In one embodiment this invention provides an isolated expression vector with a nucleic acid sequence that encodes SEQ ID NO: 2, the 26#77 protein. In further embodiments, the nucleic acid sequence is at least 80% identical to SEQ ID NO: 1. The invention also provides a host cell containing a vector that expresses a nucleic acid that encodes the 26#77 protein. Nucleic acids that encode SEQ ID NO:4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, or 28 are also provided, as well as host cells containing a vector that expresses a nucleic acid that encodes CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein.

[0015] In one embodiment this invention provides a method for determining the presence or absence of a colorectal cancer cell in a patient, by determining the level of a target protein, the 26#77 protein including the sequence shown in SEQ ID NO: 2. Levels of the 26#77 protein are determined in a biological sample from the patient, thereby determining the presence or absence of the colorectal cancer cell in the patient. In one aspect the 26#77 protein levels are determined by using an antibody specific for the 26#77 protein. The antibody can be a polyclonal antibody or a monoclonal antibody. In a further aspect, the antibody can be labeled and the label can be a fluorescent label.

[0016] In one embodiment this invention provides a method for determining the presence or absence of a colorectal cancer cell in a patient, by determining the level of a target protein, the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein including the sequences shown in SEQ ID NO:4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, or 28. Levels of the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein are determined in a biological sample from the patient, thereby determining the presence or absence of the colorectal cancer cell in the patient. In one aspect the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein levels are determined by using an antibody specific for the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein. The antibody can be a polyclonal antibody or a monoclonal antibody. In a further aspect, the antibody can be labeled and the label can be a fluorescent label.

[0017] In a further embodiment, the step of determination of the level of the 26#77 protein is carried out by comparing the amount of the 26#77 protein in the biological sample to the

amount of the 26#77 protein in a reference sample. In one aspect, the reference sample is from normal colorectal tissue. In another aspect, the determination of 26#77 protein level is made when the patient is undergoing a therapeutic regimen to treat colorectal cancer. In a further aspect, the determination of 26#77 protein level is made when the patient is suspected of having colorectal cancer. Similarly, the step of determination of the level of the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein is carried out by comparing the amount of the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein in the biological sample to the amount of the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein in a reference sample. In one aspect, the reference sample is from normal colorectal tissue. In another aspect, the determination of CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein level is made when the patient is undergoing a therapeutic regimen to treat colorectal cancer. In a further aspect, the determination of CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein level is made when the patient is suspected of having colorectal cancer.

[0018] In one embodiment, the present invention provides a method for treating a cancer that overexpresses a 26#77 gene product by administering a therapeutically effective amount of an inhibitor of 26#77 gene product to a patient who has a cancer that overexpresses 26#77. The inhibitor of the 26#77 gene product can be an antibody, an antisense RNA molecule, or an inhibitory RNA molecule. Similarly, the present invention provides a method for treating a cancer that overexpresses a CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene product by administering a therapeutically effective amount of an inhibitor of CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene product to a patient who has a cancer that overexpresses CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188. The inhibitor of the CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene product can be an antibody, an antisense RNA molecule, or an inhibitory RNA molecule.

DEFINITIONS

[0019] The phrase "determining the level of a target nucleic acid " refers to any method that can be used to detect increased copy number of a genomic sequence or increased expression level of a target gene. Methods for determining increased copy number are well known and include nucleic acid hybridization methods as described below. Methods for determining the level of expression of a particular gene are well known in the art. Such methods include RT-PCR, real-time PCR, use of antibodies against the gene products, and the like. As explained below, methods of the invention are used to detect increased copy number or overexpression of a gene referred to here as 26#77 or to detect overexpression CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 genes. Typically, overexpression of a particular gene is at least about 2 times, usually at least about 5 times the level of expression in a normal cell from the same tissue.

[0020] The terms "26#77 protein" or "26#77 polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 1 or SEQ ID NO: 2. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 1 or SEQ ID NO: 2, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0021] The terms "copine 1(CPNE 1) protein" or "copine 1(CPNE1) polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 3 or SEQ ID NO: 4. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 3 or SEQ ID NO: 4, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0022] The terms "integrin B4 binding protein (ITGB4BP) protein" or "integrin B4 binding protein (ITGB4BP) polynucleotide" or refer to nucleic acid and polypeptide polymorphic

variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 5 or SEQ ID NO: 6. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 5 or SEQ ID NO: 6, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0023] The terms "RNA export homolog (RAE) protein" or "RNA export homolog (RAE) polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 7 or SEQ ID NO: 8. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 7 or SEQ ID NO: 8, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0024] The terms "bone morphogenic protein 7 (BMP7) protein" or "bone morphogenic protein 7 (BMP7) polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 9 or SEQ ID NO: 10. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 9 or SEQ ID NO: 10, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0025] The terms "G protein, alpha stimulating activity polypeptide 1 (GNAS) protein" or "G protein, alpha stimulating activity polypeptide 1 (GNAS) polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 11 or SEQ ID NO: 12. Typically such genes or proteins have a

sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 11 or SEQ ID NO: 12, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not
5 limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0026] The terms "eukaryotic translation initiation factor 2, subunit 2 beta (EIF2S2) protein" or "eukaryotic translation initiation factor 2, subunit 2 beta (EIF2S2)

polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles,

10 mutants, and interspecies homologues of SEQ ID NO: 13 or SEQ ID NO: 14. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 13 or SEQ ID NO: 14, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a
15 mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0027] The terms "dynein light chain A2 (DNCL2A) protein" or "dynein light chain A2 (DNCL2A) polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants,

20 alleles, mutants, and interspecies homologues of SEQ ID NO: 15 or SEQ ID NO: 16.

Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 15 or SEQ ID NO: 16, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically
25 from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0028] The terms "proteasome subunit α -7 (PSMA7) protein" or "proteasome subunit α -7 (PSMA7) polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants,

30 alleles, mutants, and interspecies homologues of SEQ ID NO: 17 or SEQ ID NO: 18.

Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID

NO: 17 or SEQ ID NO: 18, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0029] The terms "activity dependent neuroprotector (ADNP) protein" or "activity dependent neuroprotector (ADNP) polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 19 or SEQ ID NO: 20. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 19 or SEQ ID NO: 20, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0030] The terms "C20orf129 protein" or "C20orf129 polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 21 or SEQ ID NO: 22. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 21 or SEQ ID NO: 22, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0031] The terms "C20orf52 protein" or "C20orf52 polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 23 or SEQ ID NO: 24. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 23 or SEQ ID NO: 24, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not

limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0032] The terms "C20orf20 protein" or "C20orf20 polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 25 or SEQ ID NO: 26. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 25 or SEQ ID NO: 26, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0033] The terms "C20orf188 protein" or "C20orf188 polynucleotide" or refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologues of SEQ ID NO: 27 or SEQ ID NO: 28. Typically such genes or proteins have a sequence that has greater than about 70% nucleotide sequence identity, usually 80%, 85%, 90% or 99% or greater sequence identity to SEQ ID NO: 27 or SEQ ID NO: 28, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more residues. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or other mammal. These terms include both naturally occurring or recombinant forms.

[0034] A "biological sample" as used herein is a sample of biological tissue or fluid that contains nucleic acids or polypeptides, e.g., of a 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein, polynucleotide or transcript. Such samples include, but are not limited to, tissue isolated from humans, or rodents, e.g., mice, and rats. Biological samples may also include sections of tissues such as biopsy and autopsy samples, frozen sections taken for histologic purposes, blood, plasma, serum, sputum, stool, tears, mucus, hair, skin, etc. Biological samples also include explants and primary and/or transformed cell cultures derived from patient tissues. A biological sample is typically obtained from a eukaryotic organism, most preferably a mammal such as a primate e.g., chimpanzee or human; cow; dog; cat; a rodent, e.g., guinea pig, rat, mouse; rabbit; or a bird; reptile; or fish. Livestock and domestic animals are of particular interest.

[0035] "Providing a biological sample" means to obtain a biological sample for use in methods described in this invention. Most often, this will be done by removing a sample of cells from an animal, but can also be accomplished by using previously isolated cells (e.g., isolated by another person, at another time, and/or for another purpose), or by performing the methods of the invention in vivo. Archival tissues, having treatment or outcome history, will be particularly useful.

[0036] The terms "identical" or percent "identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (e.g., greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%, or higher identity over a specified region, when compared and aligned for maximum correspondence over a comparison window or designated region) as measured using a BLAST or BLAST 2.0 sequence comparison algorithms with default parameters described below, or by manual alignment and visual inspection. Such sequences are then said to be "substantially identical." This definition also refers to, or may be applied to, the complement of a test sequence. The definition also includes sequences that have deletions and/or additions, as well as those that have substitutions, as well as naturally occurring, e.g., polymorphic or allelic variants, and man-made variants. As described below, the preferred algorithms can account for gaps and the like. Preferably, identity exists over a region that is at least about 25 amino acids or nucleotides in length, or more preferably over a region that is 50-100 amino acids or nucleotides in length.

[0037] For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Preferably, default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

[0038] A "comparison window", as used herein, includes reference to a segment of one of the number of contiguous positions selected from the group consisting typically of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence

may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math.

5 2:482-489, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443-453, by the search for similarity method of Pearson and Lipman (1988) Proc. Nat'l. Acad. Sci. USA 85:2444-2448, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and
10 visual inspection (see, e.g., Ausubel, et al. (eds. 1995 and supplements) Current Protocols in Molecular Biology Lippincott.

[0039] The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul (1993) Proc. Nat'l Acad. Sci. USA 90:5873-5887). One measure of similarity provided by the BLAST algorithm is the smallest sum
15 probability ($P(N)$), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001. Log values may be large
20 negative numbers, e.g., 5, 10, 20, 30, 40, 40, 70, 90, 110, 150, 170, etc.

[0040] An indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second
25 polypeptide, e.g., where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below. Yet another indication that two nucleic acid sequences are substantially identical is that the same primers can be used to amplify the sequences.

30 [0041] A "host cell" is a naturally occurring cell or a transformed cell that contains an expression vector and supports the replication or expression of the expression vector. Host cells may be cultured cells, explants, cells in vivo, and the like. Host cells may be

prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells, such as CHO, HeLa, and the like (see, e.g., the American Type Culture Collection catalog or web site).

[0042] The terms "isolated," "purified," or "biologically pure" refer to material that is substantially or essentially free from components that normally accompany it as found in its native state. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein or nucleic acid that is the predominant species present in a preparation is substantially purified. In particular, an isolated nucleic acid is separated from some open reading frames that naturally flank the gene and encode proteins other than protein encoded by the gene. The term "purified" in some embodiments denotes that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. Preferably, it means that the nucleic acid or protein is at least 85% pure, more preferably at least 95% pure, and most preferably at least 99% pure. "Purify" or "purification" in other embodiments means removing at least one contaminant from the composition to be purified. In this sense, purification does not require that the purified compound be homogenous, e.g., 100% pure.

[0043] The terms "polypeptide," "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers, those containing modified residues, and non-naturally occurring amino acid polymers.

[0044] The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function similarly to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, e.g., an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs may have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to

chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions similarly to a naturally occurring amino acid.

[0045] Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical

5 Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

[0046] "Conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino
10 acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical or associated, e.g., naturally contiguous, sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode most proteins. For instance, the codons GCA, GCC, GCG, and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can
15 be altered to another of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes silent variations of the nucleic acid. In certain contexts each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and
20 TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, a silent variation of a nucleic acid which encodes a polypeptide is implicit in a described sequence with respect to the expression product, but not necessarily with respect to actual probe sequences.

[0047] As to amino acid sequences, one of skill will recognize that individual substitutions,
25 deletions, or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds, or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such
30 conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention. Typically conservative substitutions for one another: 1) Alanine (A), Glycine (G); 2) Aspartic acid (D), Glutamic acid (E); 3)

Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); 7) Serine (S), Threonine (T); and 8) Cysteine (C), Methionine (M) (see, e.g., Creighton (1984) Proteins Freeman).

5 [0048] "Nucleic acid" or "oligonucleotide" or "polynucleotide" or grammatical equivalents used herein means at least two nucleotides covalently linked together. Oligonucleotides are typically less than about 100 nucleotides in length. A nucleic acid of the present invention will generally contain phosphodiester bonds, although in some cases, nucleic acid analogs are included that may have at least one different linkage, e.g.,
10 phosphoramidate, phosphorothioate, phosphorodithioate, or O-methylphosphoroamidite linkages (see Eckstein (1992) *Oligonucleotides and Analogues: A Practical Approach* Oxford University Press); and peptide nucleic acid backbones and linkages. For example, peptide nucleic acids (PNA) which includes peptide nucleic acid analogs can be used in the invention.

15 [0049] The nucleic acids may be single stranded or double stranded, as specified, or contain portions of both double stranded or single stranded sequence. As will be appreciated by those in the art, the depiction of a single strand also defines the sequence of the complementary strand; thus the sequences described herein also provide the complement of the sequence. The nucleic acid may be DNA, both genomic and cDNA, RNA, or a hybrid, where the
20 nucleic acid may contain combinations of deoxyribo- and ribo-nucleotides, and combinations of bases, including uracil, adenine, thymine, cytosine, guanine, inosine, xanthine hypoxanthine, isocytosine, isoguanine, etc. "Transcript" typically refers to a naturally occurring RNA, e.g., a pre-mRNA, hnRNA, or mRNA. As used herein, the term "nucleoside" includes nucleotides and nucleoside and nucleotide analogs, and modified
25 nucleosides such as amino modified nucleosides. In addition, "nucleoside" includes non-naturally occurring analog structures. Thus, e.g., the individual units of a peptide nucleic acid, each containing a base, are referred to herein as a nucleoside.

[0050] A "label" or a "detectable moiety" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, chemical, or other physical means. For
30 example, useful labels include ^{32}P , fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins or other entities which can be made detectable, e.g., by incorporating a radiolabel into the peptide or used to

detect antibodies specifically reactive with the peptide. The labels may be incorporated into the ovarian cancer nucleic acids, proteins and antibodies at any position. Any method known in the art for conjugating the antibody to the label may be employed, including those methods described by Hunter, et al. (1962) *Nature* 194:495-496; David, et al. (1974) *Biochemistry* 13:1014-1021; Pain, et al. (1981) *J. Immunol. Meth.* 40:219-230; and Nygren (1982) *J. Histochem. and Cytochem.* 30:407-412.

[0051] An "effector" or "effector moiety" or "effector component" is a molecule that is bound (or linked, or conjugated), either covalently, through a linker or a chemical bond, or non-covalently, through ionic, van der Waals, electrostatic, or hydrogen bonds, to an antibody. The "effector" can be a variety of molecules including, e.g., detection moieties including radioactive compounds, fluorescent compounds, an enzyme or substrate, tags such as epitope tags, a toxin; activatable moieties, a chemotherapeutic agent; a lipase; an antibiotic; or a radioisotope emitting "hard" e.g., beta radiation.

[0052] A "labeled nucleic acid probe or oligonucleotide" is one that is bound, either covalently, through a linker or a chemical bond, or non-covalently, through ionic, van der Waals, electrostatic, or hydrogen bonds to a label such that the presence of the probe may be detected by detecting the presence of the label bound to the probe. Alternatively, method using high affinity interactions may achieve the same results where one of a pair of binding partners binds to the other, e.g., biotin, streptavidin.

[0053] The term "probe" or a "nucleic acid probe", as used herein, is defined to be a collection of one or more nucleic acid fragments whose hybridization to a sample can be detected. The probe may be unlabeled or labeled as described below so that its binding to the target or sample can be detected. Particularly in the case of arrays, either probe or target nucleic acids may be affixed to the array. Whether the array comprises "probe" or "target" nucleic acids will be evident from the context. Similarly, depending on context, either the probe, the target, or both can be labeled. In some embodiments, the probe may be a member of an array of spotted nucleic acids. Techniques capable of producing high density arrays can also be used for this purpose (see, e.g., Fodor (1991) *Science* 767-773; Johnston (1998) *Curr. Biol.* 8: R171-R174; Schummer (1997) *Biotechniques* 23: 1087-1092; Kern (1997) *Biotechniques* 23: 120-124; U.S. Patent No. 5,143,854). One of skill will recognize that the precise sequence of the particular probes described herein can be modified to a certain degree to produce probes that are "substantially identical" to the disclosed probes, but retain the

ability to specifically bind to (i.e., hybridize specifically to) the same targets or samples as the probe from which they were derived. In addition, those of skill will recognize that a probe can specifically bind to all or a fragment of a target nucleic acid. Such modifications are specifically covered by reference to the individual probes described herein.

5 [0054] The term "recombinant" when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, e.g., recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or
10 express native genes that are otherwise abnormally expressed, under expressed or not expressed at all. By the term "recombinant nucleic acid" herein is meant nucleic acid, originally formed in vitro, in general, by the manipulation of nucleic acid, e.g., using polymerases and endonucleases, in a form not normally found in nature. In this manner, operably linkage of different sequences is achieved. Thus an isolated nucleic acid, in a linear
15 form, or an expression vector formed in vitro by ligating DNA molecules that are not normally joined, are both considered recombinant for the purposes of this invention. It is understood that once a recombinant nucleic acid is made and reintroduced into a host cell or organism, it will replicate non-recombinantly, e.g., using the in vivo cellular machinery of the host cell rather than in vitro manipulations; however, such nucleic acids, once produced
20 recombinantly, although subsequently replicated non-recombinantly, are still considered recombinant for the purposes of the invention. Similarly, a "recombinant protein" is a protein made using recombinant techniques, e.g., through the expression of a recombinant nucleic acid as depicted above.

[0055] The term "heterologous" when used with reference to portions of a nucleic acid
25 indicates that the nucleic acid comprises two or more subsequences that are not normally found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences, e.g., from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein will often refer to two
30 or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

[0056] A "promoter" is defined as an array of nucleic acid control sequences that direct transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A "constitutive" promoter is a promoter that is active under most environmental and developmental conditions. An "inducible" promoter is a promoter that is active under environmental or developmental regulation. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence (such as a promoter, or array of transcription factor binding sites) and a second nucleic acid sequence, e.g., wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

[0057] An "expression vector" is a nucleic acid construct, generated recombinantly or synthetically, with a series of specified nucleic acid elements that permit transcription of a particular nucleic acid in a host cell. The expression vector can be part of a plasmid, virus, or nucleic acid fragment. Typically, the expression vector includes a nucleic acid to be transcribed operably linked to a promoter.

[0058] The phrase "selectively (or specifically) hybridizes to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent hybridization conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

[0059] The phrase "stringent hybridization conditions" refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acids, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in "Overview of principles of hybridization and the strategy of nucleic acid assays" in Tijssen (1993) *Hybridization with Nucleic Probes (Laboratory Techniques in Biochemistry and Molecular Biology)* (vol. 24) Elsevier. Generally, stringent conditions are selected to be about 5-10° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic acid concentration) at which 50% of the probes complementary to the target hybridize to the target

sequence at equilibrium (as the target sequences are present in excess, at T_m , 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is typically at least two times background, preferably 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, 5x SSC, and 1% SDS, incubating at 42° C, or, 5x SSC, 1% SDS, incubating at 65° C, with wash in 0.2x SSC, and 0.1% SDS at 65° C. For PCR, a temperature of about 36° C is typical for low stringency amplification, although annealing temperatures may vary between about 32-48° C depending on primer length. For high stringency PCR amplification, a temperature of about 62° C is typical, although high stringency annealing temperatures can range from about 50° C to about 65° C, depending on the primer length and specificity. Typical cycle conditions for both high and low stringency amplifications include a denaturation phase of 90-95° C for 30-120 sec, an annealing phase lasting 30-120 sec, and an extension phase of about 72° C for 1-2 min. Protocols and guidelines for low and high stringency amplification reactions are available, e.g., in Innis, et al. (1990) *PCR Protocols: A Guide to Methods and Applications* Academic Press, N.Y.

[0060] "Inhibitors", and "modulators" of polynucleotides of the invention are used to refer to molecules or agents that inhibit or modulate the oncogenic effects of the proteins described here. Such agents can be identified using in vitro and in vivo assays described below. Inhibitors are compounds that, e.g., bind to, partially or totally block activity, decrease, prevent, delay activation, inactivate, desensitize, or down regulate the activity or expression of the 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 proteins described here. Such agents include, for example, antisense or inhibitory RNAs which inhibit expression of the 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene. Inhibitors also include antibodies that bind specifically to 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 proteins. In some embodiments, humanized antibodies are used as inhibitors, and can be used therapeutically.

Assays for inhibitors include, *e.g.*, expressing the 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein *in vitro*, in cells, or cell membranes, applying test compounds, and then determining the functional effects on activity (*e.g.*, changes in growth of the cell). Changes in cell growth could be any property associated with a neoplastic phenotype, for example, cell viability, formation of foci, anchorage independence, semi-solid or soft agar growth, change in contact inhibition or density limitation of growth, loss of growth factor or serum requirements, change in cell morphology, gain or loss of immortalization, gain or loss of tumor specific markers, ability to form or suppress tumors when injected into suitable animal hosts, and/or immortalization of the cell.

[0061] "Tumor cell" refers to pre-cancerous, cancerous, and normal cells in a tumor.

[0062] "Cancer cells," "transformed" cells or "transformation" in tissue culture, refers to spontaneous or induced phenotypic changes that do not necessarily involve the uptake of new genetic material. Although transformation can arise from infection with a transforming virus and incorporation of new genomic DNA, or uptake of exogenous DNA, it can also arise spontaneously or following exposure to a carcinogen, thereby mutating an endogenous gene. Transformation is typically associated with phenotypic changes, such as immortalization of cells, aberrant growth control, non-morphological changes, and/or malignancy.

[0063] As used herein, "antibody" includes reference to an immunoglobulin molecule immunologically reactive with a particular antigen, and includes both polyclonal and monoclonal antibodies. The term also includes genetically engineered forms such as chimeric antibodies (*e.g.*, humanized murine antibodies) and heteroconjugate antibodies (*e.g.*, bispecific antibodies). The term "antibody" also includes antigen binding forms of antibodies, including fragments with antigen-binding capability (*e.g.*, Fab', F(ab')₂, Fab, Fv and rIgG. *See also*, Pierce Catalog and Handbook, 1994-1995 (Pierce Chemical Co., Rockford, IL). *See also*, *e.g.*, Kuby, J., *Immunology*, 3rd Ed., W.H. Freeman & Co., New York (1998). The term also refers to recombinant single chain Fv fragments (scFv). The term antibody also includes bivalent or bispecific molecules, diabodies, triabodies, and tetrabodies. Bivalent and bispecific molecules are described in, *e.g.*, Kostelny *et al.* (1992) *J Immunol* 148:1547, Pack and Pluckthun (1992) *Biochemistry* 31:1579, Hollinger *et al.*, 1993, *supra*, Gruber *et al.* (1994) *J Immunol* :5368, Zhu *et al.* (1997) *Protein Sci* 6:781, Hu *et al.*

(1996) *Cancer Res.* 56:3055, Adams *et al.* (1993) *Cancer Res.* 53:4026, and McCartney, *et al.* (1995) *Protein Eng.* 8:301.

[0064] An antibody immunologically reactive with a particular antigen can be generated by recombinant methods such as selection of libraries of recombinant antibodies in phage or similar vectors, *see, e.g.*, Huse *et al.*, *Science* 246:1275-1281 (1989); Ward *et al.*, *Nature* 341:544-546 (1989); and Vaughan *et al.*, *Nature Biotech.* 14:309-314 (1996), or by immunizing an animal with the antigen or with DNA encoding the antigen.

[0065] Typically, an immunoglobulin has a heavy and light chain. Each heavy and light chain contains a constant region and a variable region, (the regions are also known as "domains"). Light and heavy chain variable regions contain four "framework" regions interrupted by three hypervariable regions, also called "complementarity-determining regions" or "CDRs". The extent of the framework regions and CDRs have been defined. The sequences of the framework regions of different light or heavy chains are relatively conserved within a species. The framework region of an antibody, that is the combined framework regions of the constituent light and heavy chains, serves to position and align the CDRs in three dimensional space.

[0066] The CDRs are primarily responsible for binding to an epitope of an antigen. The CDRs of each chain are typically referred to as CDR1, CDR2, and CDR3, numbered sequentially starting from the N-terminus, and are also typically identified by the chain in which the particular CDR is located. Thus, a V_H CDR3 is located in the variable domain of the heavy chain of the antibody in which it is found, whereas a V_L CDR1 is the CDR1 from the variable domain of the light chain of the antibody in which it is found.

[0067] References to " V_H " or a " VH " refer to the variable region of an immunoglobulin heavy chain of an antibody, including the heavy chain of an Fv, scFv, or Fab. References to " V_L " or a " VL " refer to the variable region of an immunoglobulin light chain, including the light chain of an Fv, scFv, dsFv or Fab.

[0068] The phrase "single chain Fv" or "scFv" refers to an antibody in which the variable domains of the heavy chain and of the light chain of a traditional two chain antibody have been joined to form one chain. Typically, a linker peptide is inserted between the two chains to allow for proper folding and creation of an active binding site.

[0069] A "chimeric antibody" is an immunoglobulin molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, e.g., an enzyme, toxin, hormone, growth factor, drug, etc.; or (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity.

[0070] A "humanized antibody" is an immunoglobulin molecule which contains minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework (FR) regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin (Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992)). Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-327 (1988); Verhoeven *et al.*, *Science* 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such humanized antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species.

[0071] "Epitope" or "antigenic determinant" refers to a site on an antigen to which an antibody binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes

formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance.

5 See, e.g., Epitope Mapping Protocols in Methods in Molecular Biology, Vol. 66, Glenn E. Morris, Ed (1996).

BRIEF DESCRIPTION OF THE DRAWINGS

[0072] Figure 1 illustrates a comparison of Northern and Western blots showing that both RNA and protein expression of 26#77 are higher in colorectal cancers with an amplified
10 26#77 gene (T) compared to the patients' normal colorectal tissue (N).

[0073] Figure 2 provides gene names, symbols, Unigene ID numbers, and accession numbers of reference DNA sequences for CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, and C20orf188.

DETAILED DESCRIPTION

15 [0074] This invention provides novel therapeutic and diagnostic methods for treatment and detection of cancer, as well as methods for screening for compositions which can be used to treat cancer. As shown below, the invention is based, at least in part, on the discovery that 26#77 is overexpressed in colorectal and breast cancer cells; and that CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20,
20 or C20orf188 are overexpressed in colorectal cancer cells. The overexpression of these genes therefore facilitates progression of carcinogenesis.

METHODS OF SCREENING FOR INCREASED COPY NUMBER OR OVEREXPRESSION OF GENES

[0075] In one aspect, 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2,
25 DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 genes (or their expression levels) are detected in different patient samples for which either diagnosis or prognosis information is desired. For example, the presence of cancer is evaluated by a determination of the increased copy number of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188
30 genes in the patient. Methods of evaluating the presence and/or copy number of a particular

gene or to determine the presence or absence of polymorphisms in the gene are well known to those of skill in the art. For example, hybridization based assays can be used for these purposes.

Hybridization-based assays

5 [0076] Hybridization assays can be used to detect copy number of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 function. Hybridization-based assays include, but are not limited to, traditional "direct probe" methods such as Southern blots or *in situ* hybridization (e.g., FISH), and "comparative probe" methods such as comparative genomic hybridization
10 (CGH). The methods can be used in a wide variety of formats including, but not limited to substrate- (e.g. membrane or glass) bound methods or array-based approaches as described below.

[0077] In a typical *in situ* hybridization assay, cells or tissue sections are fixed to a solid support, typically a glass slide. If a nucleic acid is to be probed, the cells are typically
15 denatured with heat or alkali. The cells are then contacted with a hybridization solution at a moderate temperature to permit annealing of labeled probes specific to the nucleic acid sequence encoding the protein. The targets (e.g., cells) are then typically washed at a predetermined stringency or at an increasing stringency until an appropriate signal to noise ratio is obtained.

20 [0078] The probes are typically labeled, e.g., with radioisotopes or fluorescent reporters. Preferred probes are sufficiently long so as to specifically hybridize with the target nucleic acid(s) under stringent conditions. The preferred size range is from about 200 bp to about 1000 bases.

[0079] In some applications it is necessary to block the hybridization capacity of repetitive
25 sequences. Thus, in some embodiments, tRNA, human genomic DNA, or Cot-1 DNA is used to block non-specific hybridization.

[0080] In comparative genomic hybridization methods a first collection of (sample) nucleic acids (e.g. from a possible tumor) is labeled with a first label, while a second collection of (control) nucleic acids (e.g. from a healthy cell/tissue) is labeled with a second label. The
30 ratio of hybridization of the nucleic acids is determined by the ratio of the two (first and second) labels binding to each fiber in the array. Where there are chromosomal deletions or

multiplications, differences in the ratio of the signals from the two labels will be detected and the ratio will provide a measure of the copy number.

[0081] Hybridization protocols suitable for use with the methods of the invention are described, *e.g.*, in Albertson (1984) *EMBO J.* 3: 1227-1234; Pinkel (1988) *Proc. Natl. Acad. Sci. USA* 85: 9138-9142; EPO Pub. No. 430,402; *Methods in Molecular Biology, Vol. 33: In Situ Hybridization Protocols*, Choo, ed., Humana Press, Totowa, NJ (1994), *etc.* In one particularly preferred embodiment, the hybridization protocol of Pinkel *et al.* (1998) *Nature Genetics* 20: 207-211, or of Kallioniemi (1992) *Proc. Natl Acad Sci USA* 89:5321-5325 (1992) is used.

[0082] A variety of nucleic acid hybridization formats are known to those skilled in the art. For example, common formats include sandwich assays and competition or displacement assays. Hybridization techniques are generally described in Hames and Higgins (1985) *Nucleic Acid Hybridization, A Practical Approach*, IRL Press; Gall and Pardue (1969) *Proc. Natl. Acad. Sci. USA* 63: 378-383; and John *et al.* (1969) *Nature* 223: 582-587.

[0083] The sensitivity of the hybridization assays may be enhanced through use of a nucleic acid amplification system that multiplies the target nucleic acid being detected. Examples of such systems include the polymerase chain reaction (PCR) system and the ligase chain reaction (LCR) system. Other methods recently described in the art are the nucleic acid sequence based amplification (NASBAO, Cangene, Mississauga, Ontario) and Q Beta Replicase systems.

[0084] Typically, labeled signal nucleic acids are used to detect hybridization. The labels may be incorporated by any of a number of means well known to those of skill in the art. Means of attaching labels to nucleic acids include, for example nick translation, or end-labeling by kinasing of the nucleic acid and subsequent attachment (ligation) of a linker joining the sample nucleic acid to a label (*e.g.*, a fluorophore). A wide variety of linkers for the attachment of labels to nucleic acids are also known. In addition, intercalating dyes and fluorescent nucleotides can also be used.

[0085] Detectable labels suitable for use in the present invention include any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present invention include biotin for staining with labeled streptavidin conjugate, magnetic beads (*e.g.*, DynabeadsTM), fluorescent labels (*e.g.*, fluorescein, texas red, rhodamine, green fluorescent protein, and the like, *see, e.g.*, Molecular

Probes, Eugene, Oregon, USA), radiolabels (e.g., ^3H , ^{125}I , ^{35}S , ^{14}C , or ^{32}P), enzymes (e.g., horse radish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold (e.g., gold particles in the 40 -80 nm diameter size range scatter green light with high efficiency) or colored glass or plastic (e.g., polystyrene, polypropylene, latex, etc.) beads. Patents teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241.

[0086] The label may be added to the nucleic acids prior to, or after the hybridization. So called "direct labels" are detectable labels that are directly attached to or incorporated into the sample or probe nucleic acids prior to hybridization. In contrast, so called "indirect labels" are joined to the hybrid duplex after hybridization. Often, the indirect label is attached to a binding moiety that has been attached to the target nucleic acid prior to the hybridization. Thus, for example, the target nucleic acid may be biotinylated before the hybridization. After hybridization, an avidin-conjugated fluorophore will bind the biotin bearing hybrid duplexes providing a label that is easily detected. For a detailed review of methods of labeling nucleic acids and detecting labeled hybridized nucleic acids see *Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 24: Hybridization With Nucleic Acid Probes*, P. Tijssen, ed. Elsevier, N.Y., (1993)).

[0087] The methods of this invention are particularly well suited to array-based hybridization formats. For a description of one preferred array-based hybridization system see Pinkel *et al.* (1998) *Nature Genetics*, 20: 207-211.

[0088] Arrays are a multiplicity of different "probe" or "target" nucleic acids (or other compounds) attached to one or more surfaces (e.g., solid, membrane, or gel). In a preferred embodiment, the multiplicity of nucleic acids (or other moieties) is attached to a single contiguous surface or to a multiplicity of surfaces juxtaposed to each other.

[0089] In an array format a large number of different hybridization reactions can be run essentially "in parallel." This provides rapid, essentially simultaneous, evaluation of a number of hybridizations in a single "experiment". Methods of performing hybridization reactions in array based formats are well known to those of skill in the art (see, e.g., Pastinen (1997) *Genome Res.* 7: 606-614; Jackson (1996) *Nature Biotechnology* 14:1685; Chee (1995) *Science* 274: 610; WO 96/17958, Pinkel *et al.* (1998) *Nature Genetics* 20: 207-211).

[0090] Arrays, particularly nucleic acid arrays can be produced according to a wide variety of methods well known to those of skill in the art. For example, in a simple embodiment,

"low density" arrays can simply be produced by spotting (*e.g.* by hand using a pipette) different nucleic acids at different locations on a solid support (*e.g.* a glass surface, a membrane, *etc.*).

[0091] The DNA used to prepare the arrays of the invention is not critical. For example the arrays can include genomic DNA, *e.g.* overlapping clones that provide a high resolution scan of a portion of the genome containing the desired gene, or of the gene itself. Genomic nucleic acids can be obtained from, *e.g.*, HACs, MACs, YACs, BACs, PACs, P1s, cosmids, plasmids, inter-Alu PCR products of genomic clones, restriction digests of genomic clones, cDNA clones, amplification (*e.g.*, PCR) products, and the like.

[0092] Arrays can also be produced using oligonucleotide synthesis technology. Thus, for example, U.S. Patent No. 5,143,854 and PCT Patent Publication Nos. WO 90/15070 and 92/10092 teach the use of light-directed combinatorial synthesis of high density oligonucleotide arrays.

Amplification-based assays.

[0093] In other embodiments, amplification-based assays can be used to measure 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene copy number in a sample. In such amplification-based assays, the nucleic acid sequences act as a template in an amplification reaction (*e.g.* Polymerase Chain Reaction (PCR)). In a quantitative amplification, the amount of amplification product will be proportional to the amount of template in the original sample. Comparison to appropriate (*e.g.* healthy tissue) controls provides a measure of the copy number.

[0094] Methods of "quantitative" amplification are well known to those of skill in the art. For example, quantitative PCR involves simultaneously co-amplifying a known quantity of a control sequence using the same primers. This provides an internal standard that may be used to calibrate the PCR reaction. Detailed protocols for quantitative PCR are provided in Innis *et al.* (1990) *PCR Protocols, A Guide to Methods and Applications*, Academic Press, Inc. N.Y.). The known nucleic acid sequence for the genes is sufficient to enable one of skill to routinely select primers to amplify any portion of the gene.

[0095] Real time PCR is another amplification technique that can be used to determine gene copy levels or levels of mRNA expression. (See, *e.g.*, Gibson *et al.*, *Genome Research* 6:995-1001, 1996; Heid *et al.*, *Genome Research* 6:986-994, 1996). Real-time PCR is a

technique that evaluates the level of PCR product accumulation during amplification. This technique permits quantitative evaluation of mRNA levels in multiple samples. For gene copy levels, total genomic DNA is isolated from a sample. For mRNA levels, mRNA is extracted from tumor and normal tissue and cDNA is prepared using standard techniques.

5 Real-time PCR can be performed, for example, using a Perkin Elmer/Applied Biosystems (Foster City, Calif.) 7700 Prism instrument. Matching primers and fluorescent probes can be designed for genes of interest using, for example, the primer express program provided by Perkin Elmer/Applied Biosystems (Foster City, Calif.). Optimal concentrations of primers and probes can be initially determined by those of ordinary skill in the art, and control (for
10 example, β -actin) primers and probes may be obtained commercially from, for example, Perkin Elmer/Applied Biosystems (Foster City, Calif.). To quantitate the amount of the specific nucleic acid of interest in a sample, a standard curve is generated using a control. Standard curves may be generated using the Ct values determined in the real-time PCR, which are related to the initial concentration of the nucleic acid of interest used in the assay.
15 Standard dilutions ranging from 10^{-10} to 10^{-6} copies of the gene of interest are generally sufficient. In addition, a standard curve is generated for the control sequence. This permits standardization of initial content of the nucleic acid of interest in a tissue sample to the amount of control for comparison purposes.

[0096] Other suitable amplification methods include, but are not limited to ligase chain
20 reaction (LCR) (*see* Wu and Wallace (1989) *Genomics* 4: 560, Landegren *et al.* (1988) *Science* 241: 1077, and Barringer *et al.* (1990) *Gene* 89: 117, transcription amplification (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86: 1173), self-sustained sequence replication (Guatelli *et al.* (1990) *Proc. Nat. Acad. Sci. USA* 87: 1874), dot PCR, and linker adapter PCR, *etc.*

25 Detection of gene expression

[0097] 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene expression level can also be assayed as a marker for cancer. In preferred embodiments, activity of the 26#77 gene is determined by a measure of gene transcript (*e.g.* mRNA), by a measure of the quantity of
30 translated protein, or by a measure of gene product activity. In additional embodiments, activity of a CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene is determined by a measure of gene

transcript (*e.g.* mRNA), by a measure of the quantity of translated protein, or by a measure of gene product activity.

[0098] Methods of detecting and/or quantifying the gene transcript (mRNA or cDNA) using nucleic acid hybridization techniques are known to those of skill in the art (see 5 *Sambrook et al. supra*). For example, one method for evaluating the presence, absence, or quantity of mRNA involves a Northern blot transfer.

[0099] The probes can be full length or less than the full length of the nucleic acid sequence encoding the protein. Shorter probes are empirically tested for specificity. Preferably nucleic acid probes are 20 bases or longer in length. (See *Sambrook et al.* for 10 methods of selecting nucleic acid probe sequences for use in nucleic acid hybridization.) Visualization of the hybridized portions allows the qualitative determination of the presence or absence of mRNA.

[0100] In another preferred embodiment, a transcript (*e.g.*, mRNA) can be measured using amplification (*e.g.* PCR) based methods as described above for directly assessing copy 15 number of DNA. In a preferred embodiment, transcript level is assessed by using reverse transcription PCR (RT-PCR). In another preferred embodiment, transcript level is assessed by using real-time PCR.

[0101] The expression level of an 26#77 gene can also be detected and/or quantified by detecting or quantifying the expressed 26#77 polypeptide. Similarly, the expression level of 20 a CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 gene can also be detected and/or quantified by detecting or quantifying the expressed CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polypeptide. The polypeptide can be detected and quantified by any of a number of means 25 well known to those of skill in the art. These may include analytic biochemical methods such as electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, and the like, or various immunological methods such as fluid or gel precipitin reactions, immunodiffusion (single or double), immunoelectrophoresis, radioimmunoassay (RIA), enzyme-linked 30 immunosorbent assays (ELISAs), immunofluorescent assays, western blotting, and the like. Immunohistochemical methods can also be used to detect 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or

C20orf188 protein. With immunohistochemical staining techniques, a cell sample is prepared, typically by dehydration and fixation, followed by reaction with labeled antibodies specific for the gene product coupled, where the labels are usually visually detectable, such as enzymatic labels, fluorescent labels, luminescent labels, and the like. A particularly sensitive staining technique suitable for use in the present invention is described by Hsu *et al.* (1980) *Am. J. Clin. Path.* 75:734-738. The isolated proteins can also be sequenced according to standard techniques to identify polymorphisms.

[0102] The 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polypeptide is detected and/or quantified using any of a number of well recognized immunological binding assays (see, e.g., U.S. Patents 4,366,241; 4,376,110; 4,517,288; and 4,837,168). For a review of the general immunoassays, see also Asai (1993) *Methods in Cell Biology Volume 37: Antibodies in Cell Biology*, Academic Press, Inc. New York; Stites & Terr (1991) *Basic and Clinical Immunology 7th Edition*.

[0103] Immunological binding assays (or immunoassays) typically utilize a "capture agent" to specifically bind to and often immobilize the analyte (polypeptide or subsequence). The capture agent is a moiety that specifically binds to the analyte. In a preferred embodiment, the capture agent is an antibody that specifically binds a polypeptide. The antibody (antipeptide) may be produced by any of a number of means well known to those of skill in the art.

[0104] Immunoassays also often utilize a labeling agent to specifically bind to and label the binding complex formed by the capture agent and the analyte. The labeling agent may itself be one of the moieties comprising the antibody/analyte complex. Thus, the labeling agent may be a labeled polypeptide or a labeled anti-antibody. Alternatively, the labeling agent may be a third moiety, such as another antibody, that specifically binds to the antibody/polypeptide complex.

[0105] In one preferred embodiment, the labeling agent is a second human antibody bearing a label. Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second can be modified with a detectable moiety, e.g., as biotin, to which a third labeled molecule can specifically bind, such as enzyme-labeled streptavidin. In some embodiments, Western blot analysis is used to detect and or quantify 26#77, CPNE

1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein.

[0106] Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G may also be used as the label agent. These proteins are normal constituents of the cell walls of streptococcal bacteria. They exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (*see, generally* Kronval, *et al.* (1973) *J. Immunol.*, 111: 1401-1406, and Akerstrom (1985) *J. Immunol.*, 135: 2589-2542).

[0107] 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein can be detected and/or quantified in cells using immunocytochemical or immunohistochemical methods. IHC (immunohistochemistry) can be performed on paraffin-embedded tumor blocks using a 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188-specific antibody. IHC is the method of colormetric or fluorescent detection of archival samples, usually paraffin-embedded, using an antibody that is placed directly on slides cut from the paraffin block. To detect and/or quantify 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 in, for example tissue culture cells or cells from a subject that are not embedded in paraffin (for example, hematopoietic cells) ICC (immunocytochemistry) can be used. ICC is like IHC but uses fresh, non-paraffin embedded cells plated onto slides and then fixed and stained.

[0108] Either polyclonal or monoclonal antibodies may be used in the immunoassays of the invention described herein. Polyclonal antibodies are preferably raised by multiple injections (*e.g.* subcutaneous or intramuscular injections) of substantially pure polypeptides or antigenic polypeptides into a suitable non-human mammal. The antigenicity of peptides can be determined by conventional techniques to determine the magnitude of the antibody response of an animal that has been immunized with the peptide. Generally, the peptides that are used to raise the anti-peptide antibodies should generally be those which induce production of high titers of antibody with relatively high affinity for the polypeptide.

[0109] Preferably, the antibodies produced will be monoclonal antibodies ("mAb's"). For preparation of monoclonal antibodies, immunization of a mouse or rat is preferred. Polyclonal antibodies can also be used.

[0110] It is also possible to evaluate an mAb to determine whether it has the same specificity as a mAb of the invention without undue experimentation by determining whether the mAb being tested prevents a mAb of the invention from binding to the subject gene product isolated as described above. If the mAb being tested competes with the mAb of the invention, as shown by a decrease in binding by the mAb of the invention, then it is likely that the two monoclonal antibodies bind to the same or a closely related epitope. Still another way to determine whether a mAb has the specificity of a mAb of the invention is to preincubate the mAb of the invention with an antigen with which it is normally reactive, and determine if the mAb being tested is inhibited in its ability to bind the antigen. If the mAb being tested is inhibited then, in all likelihood, it has the same, or a closely related, epitopic specificity as the mAb of the invention.

[0111] The assays of this invention have immediate utility in detecting/predicting the likelihood of a cancer, in estimating survival from a cancer, in screening for agents that modulate the subject gene product activity, and in screening for agents that inhibit cell proliferation.

METHODS OF SCREENING FOR GENE PRODUCT FUNCTION

[0112] Assays for 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 function can be designed to detect and/or quantify any effect that is indirectly or directly under the influence of the 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein or nucleic acid, e.g., a functional, physical, or chemical effect. Such assays can be used to test whether a biological sample comprises a functional 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein, to test whether variant 26#77 polypeptides retain function, or to identify compounds that modulate 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 activity in cells.

[0113] Typical assays useful in the present invention are those designed to test neoplastic characteristics of cancer cells. These assays include cell growth on soft agar; anchorage dependence; contact inhibition and density limitation of growth; cellular proliferation; cell death (apoptosis); cellular transformation; growth factor or serum dependence; tumor specific

marker levels; invasiveness into Matrigel; tumor growth and metastasis *in vivo*; mRNA and protein expression in cells undergoing metastasis, and other characteristics of cancer cells.

[0114] The ability of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polynucleotides to promote cell growth can also be assessed by introducing the polynucleotides into in cells and assessing the growth of those cells *in vitro* or *in vivo*.

[0115] Assays may include those designed to test the ability of test agents to bind the 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein and thereby modulate its activity.

Virtually any agent can be tested in such an assay. Such agents include, but are not limited to antibodies, natural or synthetic nucleic acids, natural or synthetic polypeptides, natural or synthetic lipids, natural or synthetic small organic molecules, and the like.

[0116] Proteins interacting with the peptide or with the protein encoded by the cDNA (e.g., 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188) can be isolated using a yeast two-hybrid system, mammalian two hybrid system, or phage display screen, etc. Targets so identified can be further used as bait in these assays to identify additional proteins that interact with 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 or are downstream of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188; which proteins are also targets for drug development (see, e.g., Fields *et al.*, *Nature* 340:245 (1989); Vasavada *et al.*, *Proc. Nat'l Acad. Sci. USA* 88:10686 (1991); Fearon *et al.*, *Proc. Nat'l Acad. Sci. USA* 89:7958 (1992); Dang *et al.*, *Mol. Cell. Biol.* 11:954 (1991); Chien *et al.*, *Proc. Nat'l Acad. Sci. USA* 9578 (1991); and U.S. Patent Nos. 5,283,173, 5,667,973, 5,468,614, 5,525,490, and 5,637,463).

[0117] Any of the assays for detecting 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 binding are amenable to high throughput screening. High throughput assays for the presence, absence, or quantification of particular nucleic acids or protein products are well known to those of skill in the art. Similarly, binding assays and reporter gene assays are similarly well known. Thus, for example, U.S. Patent 5,559,410 discloses high throughput screening methods for proteins, U.S. Patent 5,585,639 discloses high throughput screening methods for

nucleic acid binding (*i.e.*, in arrays), while U.S. Patents 5,576,220 and 5,541,061 disclose high throughput methods of screening for ligand/antibody binding.

[0118] In addition, high throughput screening systems are commercially available (*see, e.g.*, Zymark Corp., Hopkinton, MA; Air Technical Industries, Mentor, OH; Beckman Instruments, Inc. Fullerton, CA; Precision Systems, Inc., Natick, MA, *etc.*). These systems typically automate entire procedures including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. The manufacturers of such systems provide detailed protocols for various high throughput systems. Thus, for example, Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like.

RECOMBINANT PRODUCTION OF 26#77 POLYPEPTIDES

[0119] The present invention also provides methods, reagents, and vectors useful for expression of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polypeptides and nucleic acids *in vitro*. *In vitro* expression is particularly useful for production of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polypeptides.

[0120] Any number of well known host cells can be used for production of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polypeptides. Host cells may be cultured cells, cell lines, cells *in vivo*, and the like. Host cells may be prokaryotic cells such as bacterial cells, (*e.g.*, *E. coli*), or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells such as CHO, HeLa, and the like.

[0121] The particular procedure used to introduce the nucleic acids into a host cell for expression of the 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein is not critical to the invention. Any of the well known procedures for introducing foreign nucleotide sequences into host cells *in vitro* may be used. These include the use of calcium phosphate transfection, electroporation, liposome-mediated transfection, injection and microinjection, ballistic

methods, viral particles, virosomes, immunoliposomes, polycation:nucleic acid conjugates, naked DNA, artificial virions, agent-enhanced uptake of DNA, and the like.

[0122] In these embodiments of this invention, 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 nucleic acids are inserted into vectors using standard molecular biological techniques.

Vectors may be used at multiple stages of the practice of the invention, including for subcloning nucleic acids encoding components of the 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein as well as additional elements controlling protein expression, vector selectability, etc. Vectors may also be used to maintain or amplify the nucleic acids, for example by inserting the vector into prokaryotic or eukaryotic cells and growing the cells in culture. In addition, vectors may be used to introduce and express nucleic acids into cells for therapeutic or experimental purposes.

[0123] A variety of commercially or commonly available vectors and vector nucleic acids can be converted into a vector of the invention by cloning a nucleic acid encoding a 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein of the invention into the commercially or commonly available vector. A variety of common vectors suitable for this purpose are well known in the art.

[0124] In a typical embodiment, an 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polynucleotide is placed under the control of a promoter. A nucleic acid is "operably linked" to a promoter when it is placed into a functional relationship with the promoter. For instance, a promoter or enhancer is operably linked to a coding sequence if it increases or otherwise regulates the transcription of the coding sequence. Similarly, a "recombinant expression cassette" or simply an "expression cassette" is a nucleic acid construct, generated recombinantly or synthetically, with nucleic acid elements that are capable of effecting expression of a structural gene in hosts compatible with such sequences. Expression cassettes include promoters and, optionally, introns, polyadenylation signals, and transcription termination signals. Typically, the recombinant expression cassette includes a nucleic acid to be transcribed (*e.g.*, a nucleic acid encoding a desired polypeptide), and a promoter. Additional factors necessary or helpful in effecting expression may also be used as described

herein. For example, an expression cassette can also include nucleotide sequences that encode a signal sequence that directs secretion of an expressed protein from the host cell. Transcription termination signals, enhancers, and other nucleic acid sequences that influence gene expression, can also be included in an expression cassette.

- 5 [0125] An extremely wide variety of promoters are well known, and can be used in the vectors of the invention, depending on the particular application. Ordinarily, the promoter selected depends upon the cell in which the promoter is to be active. Other expression control sequences such as ribosome binding sites, transcription termination sites and the like are also optionally included. For *E. coli*, example control sequences include the T7, trp, or
10 lambda promoters, a ribosome binding site and preferably a transcription termination signal. For eukaryotic cells, the control sequences typically include a promoter which optionally includes an enhancer derived from immunoglobulin genes, SV40, cytomegalovirus, a retrovirus (*e.g.*, an LTR based promoter) etc., and a polyadenylation sequence, and may include splice donor and acceptor sequences.
- 15 [0126] For long-term, high-yield production of recombinant proteins, stable expression will often be desired. For example, cell lines which stably express a 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein can be prepared using expression vectors of the invention which contain viral origins of replication or endogenous expression elements and a selectable
20 marker gene. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth of cells which successfully express the introduced sequences in selective media. Resistant, stably transfected cells can be proliferated using tissue culture techniques appropriate to the cell
25 type. An amplification step, *e.g.*, by administration of methyldtrexate to cells transfected with a DHFR gene according to methods well known in the art, can be included.

KITS USE IN DIAGNOSTIC, RESEARCH, AND THERAPEUTIC APPLICATIONS

- [0127] For use in diagnostic, research, and therapeutic applications disclosed here, kits are also provided by the invention. In the diagnostic and research applications such kits may
30 include any or all of the following: assay reagents, buffers, 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188-specific nucleic acids or antibodies, hybridization probes and/or primers, and

the like. A therapeutic product may include sterile saline or another pharmaceutically acceptable emulsion and suspension base.

[0128] In addition, the kits may include instructional materials containing directions (*i.e.*, protocols) for the practice of the methods of this invention. While the instructional materials typically comprise written or printed materials they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this invention. Such media include, but are not limited to electronic storage media (*e.g.*, magnetic discs, tapes, cartridges, chips), optical media (*e.g.*, CD ROM), and the like. Such media may include addresses to internet sites that provide such instructional materials.

[0129] The present invention also provides for kits for screening for modulators of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188. Such kits can be prepared from readily available materials and reagents. For example, such kits can comprise one or more of the following materials: an 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 polypeptide or polynucleotide, reaction tubes, and instructions for testing the desired 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 function.

[0130] A wide variety of kits and components can be prepared according to the present invention, depending upon the intended user of the kit and the particular needs of the user. Diagnosis would typically involve evaluation of a plurality of genes or products. The genes will be selected based on correlations with important parameters in disease which may be identified in historical or outcome data.

THERAPEUTIC METHODS

Administration of Inhibitors

[0131] As noted above, inhibitors of the invention can be used to treat cancer and other diseases associated with pathological cellular proliferation. The compounds that inhibit 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 activity can be administered by a variety of methods including, but not limited to parenteral (*e.g.*, intravenous, intramuscular, intradermal, intraperitoneal, and subcutaneous routes), topical, oral, local, or transdermal administration. These methods can be used for prophylactic and/or therapeutic treatment.

The pharmaceutical compositions can be administered in a variety of unit dosage forms depending upon the method of administration. For example, unit dosage forms suitable for oral administration include powder, tablets, pills, capsules and lozenges.

5 [0132] The compositions for administration will commonly comprise an inhibitor dissolved in a pharmaceutically acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers can be used, *e.g.*, buffered saline and the like. These solutions are sterile and generally free of undesirable matter. These compositions may be sterilized by conventional, well known sterilization techniques. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as
10 pH adjusting and buffering agents, toxicity adjusting agents and the like, for example, sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate and the like. The concentration of active agent in these formulations can vary widely, and will be selected primarily based on fluid volumes, viscosities, body weight and the like in accordance with the particular mode of administration selected and the patient's needs.

15 [0133] Thus, a typical pharmaceutical composition for intravenous administration would be about 0.1 to 10 mg per patient per day. Dosages from 0.1 up to about 100 mg per patient per day may be used, particularly when the drug is administered to a secluded site and not into the blood stream, such as into a body cavity or into a lumen of an organ. Substantially higher dosages are possible in topical administration. Actual methods for preparing parenterally
20 administrable compositions will be known or apparent to those skilled in the art and are described in more detail in such publications as *Remington's Pharmaceutical Science*, 15th ed., Mack Publishing Company, Easton, Pennsylvania (1980).

[0134] The compositions containing inhibitors can be administered for therapeutic or prophylactic treatments. In therapeutic applications, compositions are administered to a
25 patient suffering from a disease (*e.g.*, colon cancer) in an amount sufficient to cure or at least partially arrest the disease and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose." Amounts effective for this use will depend upon the severity of the disease and the general state of the patient's health. Single or multiple administrations of the compositions may be administered depending on the dosage
30 and frequency as required and tolerated by the patient. In any event, the composition should provide a sufficient quantity of the agents of this invention to effectively treat the patient.

Polynucleotide inhibitors

[0135] The activity of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 protein can also be down-regulated, or entirely inhibited, by the use of antisense polynucleotides, e.g., a nucleic acid complementary to, and which can preferably hybridize specifically to, a 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 encoding mRNA. Binding of the antisense polynucleotide to the mRNA reduces the translation and/or stability of the mRNA.

[0136] Antisense polynucleotides can comprise naturally-occurring nucleotides, or synthetic species formed from naturally-occurring subunits or their close homologs. Antisense polynucleotides may also have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur containing species which are known for use in the art. Analogs are comprehended by this invention so long as they function effectively to hybridize with the ovarian cancer protein mRNA. See, e.g., Isis Pharmaceuticals, Carlsbad, CA; Sequitor, Inc., Natick, MA.

[0137] RNA interference is another mechanism to suppress gene expression in a sequence specific manner. See, e.g., Brumelkamp, et al. (2002) *Scienceexpress* (21March2002); Sharp (1999) *Genes Dev.* 13:139-141; and Cathew (2001) *Curr. Op. Cell Biol.* 13:244-248. In mammalian cells, short, e.g., 21 nt, double stranded small interfering RNAs (siRNA) have been shown to be effective at inducing an RNAi response. See, e.g., Elbashir, et al. (2001) *Nature* 411:494-498.

[0138] Ribozymes can also be used to target and inhibit transcription of 26#77, CPNE 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129, C20orf52, C20orf20, or C20orf188 nucleotide sequences. A ribozyme is an RNA molecule that catalytically cleaves other RNA molecules. Different kinds of ribozymes have been described, including group I ribozymes, hammerhead ribozymes, hairpin ribozymes, RNase P, and axhead ribozymes (see, e.g., Castanotto, et al. (1994) *Adv. Pharmacol.* 25: 289-317 for a general review of the properties of different ribozymes).

[0139] The polynucleotide inhibitors can be introduced into a cancer cell by any of a number of well known techniques. For example, the polynucleotide inhibitors can be conjugated to a binding molecule, as described in WO 91/04753. Suitable binding molecules include, but are not limited to, molecules that bind cell surface receptors on the surface of the

target cancer cell. Preferably, conjugation of the binding molecule does not substantially interfere with the ability of the binding molecule to bind to its corresponding receptor, or block entry of the inhibitory polynucleotide into the cell. Alternatively, a polynucleotide inhibitor may be introduced into a cell containing the target nucleic acid sequence by formation of an polynucleotide-lipid complex.

EXAMPLES

The following examples are offered to illustrate, but not to limit the claimed invention.

Example 1: Unknown gene 26#77 is amplified and overexpressed at the RNA and protein levels in primary human colorectal cancers.

[0140] Chromosome 20q is amplified in approximately 60% of primary human colorectal cancers. However, no definitive gene target has been identified for the amplicon in human colorectal cancers.

[0141] Unknown gene 26#77 was originally identified by virtue of its RNA expression profile in a breast cancer cell line. Recombinant 26#77 protein was expressed and used to generate antibodies specific for the 26#77 protein.

[0142] Twelve breast and colorectal cancer cell lines were tested for 26#77 DNA amplification and for RNA and protein levels. The 26#77 gene was amplified in three of twelve breast and colorectal cancer cell lines tested by Southern blot analysis or FISH. Northern blot analysis demonstrated that 26#77 RNA levels were elevated in nine of the twelve breast and colorectal cancer cell lines tested.

[0143] The 26#77 protein was predominantly localized in the nucleus. A colorectal cancer cell line (CACO2 cell line) was fractionated into cytoplasmic and nuclear fractions. Western blot analysis with the ant 26#77 polyclonal antibody demonstrated that the majority of 26#77 protein was found in the nuclear fraction. Immunocytochemical analysis of CACO2 cells also showed that 26#77 was predominantly localized in the nucleus. Similar results were obtained using a breast cancer cell line (BT474) that overexpressed 26#77.

[0144] 26#77 was also cloned into a tetracycline-inducible vector (from Invitrogen). The tet-inducible 26#77 vector was then used to transfect NIH 3T3 cells. After induction of 26#77

expression, 26#77 was localized to the cell nucleus as demonstrated by western blot analysis and immunocytochemistry.

[0145] One hundred and twenty-five primary colorectal cancers with the 20q amplicon were tested for 26#77 gene copy levels. The 26#77 gene was amplified in 60% of the 125 colorectal cancers tested by Southern blot analysis or FISH. A subset of the 125 primary colorectal cancers (40 samples total) were tested for 26#77 RNA and protein levels. Of the 26#77 amplified colorectal cancers in the subset (20 cancers total), all had elevated levels of 26#77 RNA compared to matched normal colorectal tissue as demonstrated by Northern blot analysis. Exemplary results are shown in Figure 1. Western blot analysis and immunohistochemistry demonstrated that 26#77 protein levels were also elevated in the samples. The results indicate that the 26#77 gene is a target of the 20q amplicon and is an important novel oncogene in human colorectal cancer.

Example 2: Other gene are amplified and overexpressed in primary human colorectal cancers.

[0146] Thirteen additional genes that reside on the q-arm of chromosome 20 are amplified in approximately 60% of human colorectal cancers and have concurrent upregulation of their RNA. Most genes in colorectal cancer amplicons downregulate their RNA to maintain normal levels. These thirteen genes do not, and are therefore upregulated at both the DNA and RNA level and may contribute to the cancer phenotype; *i.e.*, they may be targets of the amplification. The thirteen genes encode Copine I (CPN1), Integrin beta-4 binding protein (ITGB4BP), RNA Export I (RAE1), Bone morphogenic protein 7 (BMP7), GTP-binding protein, alpha-stimulatory (GNAS), eukaryotic translation initiation factor 2, subunit 2 (EIF2S2), Dynein light chain A2, (DNCL2A), Proteosome subunit alpha-type 7 (PSMA7), Activity dependent neuroprotector (ADNP), C20ORF129, C20ORF52, C20ORF20, and C20ORF188. Accession numbers for the genes are found in Figure 2.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

WHAT IS CLAIMED IS:

- 1 1. A method for determining the presence or absence of a colorectal
2 cancer cell in a patient, the method comprising determining the level of a target nucleic acid
3 that encodes SEQ ID NO: 2 in a biological sample from the patient, thereby determining the
4 presence or absence of the colorectal cancer cell in the patient.
- 1 2. The method of claim 1, wherein the target nucleic acid comprises a
2 sequence at least 80% identical to SEQ ID NO: 1.
- 1 3. The method of claim 1, wherein the biological sample comprises
2 isolated nucleic acids.
- 1 4. The method of claim 3, further comprising the step of amplifying
2 nucleic acids before the step of determining the level of the nucleic acid.
- 1 5. The method of claim 3, wherein the isolated nucleic acids are mRNA.
- 1 6. The method of claim 1, wherein the biological sample is colorectal
2 tissue and the step of determining the level of target nucleic acid is carried out using *in situ*
3 hybridization.
- 1 7. The method of claim 1, wherein the step of determining the level of
2 target nucleic acid is carried out using a labeled nucleic acid probe that selectively hybridizes
3 to SEQ ID NO: 1 under stringent hybridization conditions.
- 1 8. The method of claim 1, wherein the step of determining the level of
2 target nucleic acid is carried out using a nucleic acid probe immobilized to a solid support,
3 wherein the probe selectively hybridizes to SEQ ID NO: 1 under stringent hybridization
4 conditions.
- 1 9. The method of claim 1, wherein the step of determining the level of
2 target nucleic acid is carried out using Northern blot analysis.
- 1 10. The method of claim 1, wherein the step of determining the level of the
2 target nucleic acid is carried out by comparing the amount of the target nucleic acid in the
3 biological sample to the amount of the target nucleic acid in a reference sample.

- 1 11. The method of claim 10, wherein the reference sample is from normal
2 colorectal tissue.
- 1 12. The method of claim 1, wherein the patient is undergoing a therapeutic
2 regimen to treat colorectal cancer.
- 1 13. The method of claim 1, wherein the patient is suspected of having
2 colorectal cancer.
- 1 14. An isolated expression vector comprising a nucleic acid sequence that
2 encodes SEQ ID NO: 2.
- 1 15. The isolated expression vector of claim 14, wherein the nucleic acid
2 sequence is at least 80% identical to SEQ ID NO: 1.
- 1 16. A host cell comprising the expression vector of claim 14.
- 1 17. A method for determining the presence or absence of a colorectal
2 cancer cell in a patient, the method comprising determining the level of a target protein
3 comprising a sequence as shown in SEQ ID NO: 2 in a biological sample from the patient,
4 thereby determining the presence or absence of the colorectal cancer cell in the patient.
- 1 18. The method of claim 17, wherein the step of determining the level of
2 the target protein is carried out using an antibody.
- 1 19. The method of claim 18, wherein the antibody is a monoclonal
2 antibody.
- 1 20. The method of claim 18, wherein the antibody is a polyclonal
2 antibody.
- 1 21. The method of claim 18, wherein the antibody is labeled.
- 1 22 The method of claim 21, wherein the label is fluorescent.

1 23. The method of claim 17, wherein the step of determining the level of
2 the target protein is carried out by comparing the amount of the target protein in the
3 biological sample to the amount of the target protein in a reference sample.

1 24. The method of claim 23, wherein the reference sample is from normal
2 colorectal tissue.

1 25. The method of claim 17, wherein the patient is undergoing a
2 therapeutic regimen to treat colorectal cancer.

1 26. The method of claim 17, wherein the patient is suspected of having
2 colorectal cancer.

1 27. A method for treating a cancer that overexpresses a 26#77 gene
2 product comprising administering to a subject in need of such treatment a therapeutically
3 effective amount of an inhibitor of 26#77 gene product.

1 28. The method of claim 27, wherein the inhibitor of a 26#77 gene product
2 is selected from the group consisting of an antisense RNA molecule, and an inhibitory RNA
3 molecule.

1 29. A method for determining the presence or absence of a colorectal
2 cancer cell in a patient, the method comprising determining the level of a target nucleic acid
3 that encodes SEQ ID NO:4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, or 28 in a biological
4 sample from the patient, thereby determining the presence or absence of the colorectal cancer
5 cell in the patient.

1 30. A method for determining the presence or absence of a colorectal
2 cancer cell in a patient, the method comprising determining the level of a target protein
3 comprising a sequence as shown in SEQ ID NO:4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, or
4 28 in a biological sample from the patient, thereby determining the presence or absence of the
5 colorectal cancer cell in the patient.


1 31. A method for treating a cancer that overexpresses a Copine 1 (CPNE
2 1) protein, the Integrin B4 binding protein (ITGB4BP), RNA Export homolog (RAE1), bone

3 morphogenic protein 7 (BMP7), G protein, alpha stimulating activity polypeptide 1 (GNAS),
4 eukaryotic translation initiation factor 2, subunit 2 beta (EIF2S2), dynein light chain A2
5 (DNCL2A), proteosome subunit α -7 (PSMA7), activity dependent neuroprotector (ADNP),
6 C20orf129, C20orf52, C20orf20, or C20orf188 gene product comprising administering to a
7 subject in need of such treatment a therapeutically effective amount of an inhibitor of CPNE
8 1, ITGB4BP, RAE1, BMP7, GNAS, EIF2S2, DNCL2A, PSMA7, ADNP, C20orf129,
9 C20orf52, C20orf20, or C20orf188 gene product.

RNA and protein expression of 26#77 is higher in paired colon tumors vs. normals

Amplified samples Non-amplified samples

C107 C110 C114
BT474 T N T N T N T N T N T N



C107 C110 C114
BT474 T N T N T N T N

Western blot

Figure 1

INFORMAL SEQUENCE LISTING

SEQ ID NO:1

1 ctttgggggt ttgctgctgg ctctgactcc cgtcctgcga tgggttgcca cgggggaaca
 5 61 atccccaaga ggcataaact ggtgaagggg ccgaagaagg ttgagaaggt cgacaaagat
 121 gctgaattag tggcccaatg gaactattgt actctaagtc aggaatatatt aagacgacca
 181 atagttgcct gtgaacttgg cagactttat aacaaagatg ccgtcattga atttctcttg
 241 gacaaatctg cagaaaaggc tcttggggaag gcagcatctc acattaaaag cattaagaat
 301 gtgacagagc tgaagctttc tgataatcct gcctgggaag gggataaagg aaacactaaa
 10 361 ggtgacaagc acgatgacct ccagcgggag cgtttcatct gccccgttgt gggcctggag
 421 atgaacggcc gacacaggtt ctgcttcctt cgggtgctgc gctgtgtgtt ttctgagcga
 481 gccttgaaag agataaaagc ggaagtttgc cacacgtgtg gggctgcctt ccaggaggat
 541 gatgtcatcg tgctcaatgg caccaaggag gatgtggacg tgctgaagac aaggatggag
 601 gagagaaggc tgagagcgaa gctggaaaag aaaacaaaaga aaccaaggc agcagagtct
 661 gtttcaaaac cagatgtcag tgaagaagcc ccagggccat caaaagttaa gacagggaag
 15 721 cctgaagagg ccagccttga ttctagagag aagaaaacca acttggctcc caaaagcaca
 781 gcaatgaatg agagctcttc tggaaaagct gggaagcctc cgtgtggagc cacaagagg
 841 tccatcgctg acagtgaaga atcggaggcc tacaagtcct tctttaccac tcacagctcc
 901 gccaagcgct ccaaggagga gtctgcccac tgggtcacc acacgtccta ctgcttctga
 961 agcccgact gccaccgctc ctgcccaga aggttggtta gtttccactg aggcaggtcg
 20 1021 ctttgtgcct ctgagtgccg tctgtgtgtg tctctctata gttctgctc ataaagctgt
 1081 cctggccagc cttcaagctg gtgtggccac tcttgatgtg aggcgtgtcg gttccagggg
 1141 ggacatggga ggggctgcac agtggcccga ggtcatgctt gcttccacct gcaggtgcat
 1201 ttggctcctt ccatggccag gaagccctgt gggctgcact ttttatgctt gcagtagcaa
 1261 gagactccag agtcctcacc ggtgcagagt tggcacatat taattaacta aaattcta
 25 1321 gatcttgcta ccagcaataa atcaagtagg ccaagtgaag ctgggcttta aaaaggatgg
 1381 atttcaaata cactgtgccc actagaagct tcgaagggcc tcgtccctct gctacagccc
 1441 tgggaggagc caggatcctt gttggtctag ctaaatactg ttaggggagt gtgccccatc
 1501 tcatcatttc gaagatagca gagtcatagt tgggcaccg gtgattgggt tcaaaaataa
 1561 agctgggtctg cctcttctca aaaaaaaaaa aaagaaaaaa aaaaa

SEQ ID NO:2

MGCDGGTIPKRHELVKGPVKVEKVDKDAELVAQWNYCTLSQEIL
 35 RRPVACELGRLYNKDAVIEFLDKSAEKALGKAASHIKSIKNVTELKLSDNPAWEGD
 KGNTKGDKHDDLQRRARFICPVVGLMNGRHRFCFLRCCGCVFSEALKEIKAEVCHTC
 GAAFQEDDVIVLNGTKEDVDVLKTRMEERRLRKLEKTKPKAAESVSKPDVSEEAP
 GPSKVKTGKPEEASLDSREKKTNLAPKSTAMNESSGKAGKPPCGATKRSIADSESE
 40 AYKSLFTTHSSAKRSKEESAHWVTHTSYCF

SEQ ID NO: 3

Copine 1 (CPNE1) BC001142

1 ggcgaaggct ttgtagagtt cagaaatgag gctgactata aggtgctct gtgtcgtcat
 45 61 aaacagtaca tgggcaatcg ctttattcaa gtctatccaa ttactaagaa aggtatgcta
 121 gaaaagatag atatgattcg aaaaagactg cagaacttca gctatgacca gagggaaatg
 181 atactaaatc cagaggggga tgtcaactct gccaaagtct gtgcccacat aacaaatatt
 241 ccattcagca ttacaaagat ggatgttctt cagttcctag aaggaatccc agtggatgaa
 301 aatgctgtac atgttcttgt tgataacaat gggcaaggct taggacaggc attggttcag
 50 361 tttaaaaatg aagatgatgc acatggccca ctgcgtgacc ttggttcagc tgtccatttc
 421 ctgtgacct ctcattgaca aggacatcgg ctccaagtct gaccactct gcgtcctttt
 481 acaggatgtg ggagggggca gctgggctga gcttgccgg actgaacggg tgcggaaactg
 541 ctcaagccct gagttctcca agactctaca gcttgagtac cgctttgaga cagtccagaa
 601 gctacgcttt ggaatctatg acatagacaa caagacgcca gagctgaggg atgatgactt
 55 661 cctagggggt gctgagtgtt ccctaggaca gattgtgtcc agccaggtag tgactctccc
 721 cttgatgctg aagcctggaa aacctgctgg gcgggggacc atcacggtct cagctcagga

5 781 attaaaggac aatcgtgtag taaccatgga ggtagaggcc agaaacctag ataagaagga
841 cttcctggga aaatcagatc ctttcttgga gttcttccgc cagggtgatg ggaaatggca
901 cctgggtgtac agatctgagg tcatcaagaa caacctgaac cctacatgga agcggtttctc
961 agtccccgtt cagcatttct gtgggtggaa cccagcaca cccatccagg tgcaatgctc
10 1021 cgattatgac agtgacgggt cacatgatct catcgggtacc ttccacacca gcttggccca
1081 gctgcaggca gtcccggctg agtttgaatg catccaccct gagaagcagc agaaaaagaa
1141 aagctacaag aactctggaa ctatccgtgt caagatttgt cgggtagaaa cagagtactc
1201 ctttctggac tatgtgatgg gaggtgtca gatcaacttc actgtgggcg tggacttcac
1261 tggctccaat ggagaccct cctcacctga ctccctacac tacctgagtc caacaggggt
1321 caatgagtag ctgatggcac tgtggagtgt gggcagcgtg gttcaggact atgactcaga
1381 caagctgttc cctgcatttg gatttggggc ccagggttccc cctgactggc aggtctcgca
1441 tgaatttgcc ttgaatttca accccagtaa cccctactgt gcaggcatcc agggcattgt
1501 ggatgcctac cgccaagccc tgcccgaagt tcgcctctat ggccctacca actttgcacc
1561 catcatcaac catgtggcca ggtttgcagc ccaggctgca catcagggga ctgcctcgca
15 1621 atacttcatg ctggttgcgc tgactgatgg tgctgtgacg gatgtggaag ccacacgtga
1681 ggctgtgggtg cgtgcctcga acctgcccat gtcagtgatc attgtgggtg tgggtgtgtg
1741 tgactttgag gccatggagc agctggagcg tgatgggtga cccctgcata cacgttctgg
1801 gcaggctgct gcccgcgaca ttgtgcagtt tgtaccctac cgccgggttc agaagtcccc
1861 tcgggaggca ttggcacaga ccgtgctcgc agaagtggc acacaactgg tctcatactt
20 1921 cagggccag ggttggggcc cgctcaagcc acttccacc tcagccaagg atcctgcaca
1981 ggccccccag gcctagggtt ccttggggc tgtggcaagt cctcaatcct gtgtcccaga
2041 ggtccctctg ggccacaacc caacccttct cactctctc agtgctagca ctttgtattt
2101 tttgatactt ttatacttgt ttctgctttt gctgctcttg atccacctt tgctcctgac
2161 aaccctcatt caataaagac cagtgaagac caaaaaaaaa aaaaaaaaaa a

25

SEQ ID NO: 4

Copine 1 (CPNE1) BC001142

30 /translation="MAHCVTLVQLSISCDHLIDKDIGSKSDPLCVLLQDVGGGSWAEI
GRTERVRNCSSPEFSKTLQLEYRFETVQKLRFGIYDIDNKTPELRDDDFLGGAECSLG
QIVSSQVLTLPMLKPGKPAGRGTTITVSAQELKDNRVVTMEVEARNLDKKDFLGKSDP
FLEFFRQGDGKWHLVYRSEVIKNNLNPTWKRFSPVQHFCCGNPSTPIQVQCSDYDSD
GSHDLIGTFHTSLAQLQAVPAEFECIHPEKQKKKSYKNSGTIRVKICRVETEYSFLD
YVMGGCQINFVTGVDFTGSNGDPSSPDSLHYLSPTGVNEYLMALWSVGSVVQDYDSK
35 LFPAFGFGAQVPPDWQVSHEFALNFNPSNPYCAGIQGIVDAYRQALPQVRLYGPTNFA
PIINHVARFAAQAAHQGTASQYFMLLLLTGAVTDVEATREAVVRASNLPMSEVIVGV
GGADFEAMEQLDADGGPLHTRSGQAAARDIVQFVYRRFQNAPREALAQTVLAEVPTQ
LVSYFRAQGWAPLKPLPPSAKDPAQAPQA"

SEQ ID NO: 5

40 Integrin B4 bind. prot (ITGB4BP) AF047433

1 aacggaaacc tttttaggga gtccaaggta cagtcgccgc gtgaggagct tggtactggg
61 tacttggcct catggcggtc cgagcttctg tcgagaacaa ctgtgagatc ggctgctttg
121 ccaagctcac caacacctac tgtctggtag cgatcggagg ctgagagaac ttctacagt
181 tgttcgaggg cgagctctcc gataccatcc ccgtgggtgca cgcgtctatc gccggctgcc
45 241 gcatcatcgg gcgcatgtgt gtggggaaca ggcacggtct cctggtacct aacaatacca
301 ccgaccagga gctgcaacac attcgcaaca gcctcccaga cacagtgcag attaggcggg
361 tggaggagcg gctctcagcc ttgggcaatg tcaccacctg caatgactac gtggccttgg
421 tccaccaga cttggacagg gagacagaag aaattctggc agatgtgctc aagggtgaag
481 tcttcagaca gacagtggcc gaccaggtgc tagtaggaag ctactgtgtc ttcagcaatc
50 541 agggagggct ggtgcattcc aagacttcaa ttgaagacca ggatgagctg tctctcttc
601 ttcaagtccc ccttgtggcg gggactgtga accgaggcag tgagggtgatt gctgctggga
661 tgggtggtag tgactggtgt gccttctgtg gcctggacac aaccagcaca gagctgtcag
721 tgggtggagag tgtcttcaag ctgaatgaag ccagcctag caccattgcc accagcatgc
781 gggattccct cattgacagc ctcacctgag tcaccttcca agttgttcca tgggctcctg
55 841 gctctggact gtggccaacc ttctccacat tccgcccatt ctgtaccgga tgctggcagg
901 gaggtggcag agagctcact gggactgagg ggctgggcac ccaaccctt tccacctgtg

961 cttatcgctt ggatctatca ttactgcaaa aacctgctct gttgtgctgg ctggcaggcc
 1021 ctgtggctgc tggctgaggg ttctgctgtc ctgtgccacc ccattaaagt gcagttccct
 1081 ccggaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa

5 SEQ ID NO: 6

Integrin B4 bind. prot (ITGB4BP) AF047433

/translation="MAVRASFENNCEIGCFAKLTNTYCLVAIGGSENFYSVFEGELSD
 TIPVVHASIAGCRIIGRMCVGNRHGLLVPNNTTDQELQHIRNSLPDTVQIRRV EERLS
 10 ALGNVTTTCNDYVALVHPDLRETEEILADVLKVEVFRQTVADQVLVGSYCVFSNQGL
 VHPKTSIEDQDELSSLLQVPLVAGTVNRGSEVIAAGMVVNDWCAFCGLDSTSTELSVV
 ESVFKLNEAQPSTIATSMRDSLIDSLT"

SEQ ID NO: 7

RNAExportHomol. (RAE1) U84720

15 1 gcggtagtca gggcagtttc taccgcagggc ttaaggaggc ttcgggctcc tgggatttct
 61 gtccgcgctc ctggccctcg tccctcgccg cagagcaggt tgcgaaactc ctcagaccct
 121 tctgtcccg gccgcgctt tccgcggggg cgagaccccc aggttcaaaa tgagcctgtt
 181 tggaacaacc tcaggttttg gaaccagtg gaccagcatg tttggcagtg caactacaga
 241 caatcacaat cccatgaagg atattgaagt aacatcatct cctgatgata gcattggttg
 20 301 tctgtctttt agcccaccaa ccttgccggg gaactttctt attgcaggat catgggctaa
 361 tgatgttcgc tgcctgggaag ttcaagacag tggacagacc attccaaaag cccagcagat
 421 gcacactggg cctgtgcttg atgtctgctg gagtgcagat gggagcaaaag tgtttacggc
 481 atcgtgtgat aaaactgcca aaatgtggga cctcagcagt aaccaagcga tacagatcgc
 541 acagcatgat gctcctgtta aaacatcca ttggatcaaa gctccaaact acagctgtgt
 25 601 gatgactggg agctgggata agactttaaa gttttgggat actcgatcgt caaatcctat
 661 gatggttttg caactccctg aaagggtgta ctgtgctgac gtgatatacc ccatggctgt
 721 ggtggcaact gcagagaggg gcctgattgt ctatcagcta gagaatcaac cttctgaatt
 781 caggaggata gaatctccac tgaaacatca gcacgggtgt gtggctattt ttaaagacaa
 841 acagaacaag cctactggtt ttgccctggg aagtatcgag gggagagttg ctattcacta
 30 901 tatcaacccc ccgaaccccg ccaaagataa cttcacctt aaatgtcatc gatctaattg
 961 aaccaacact tcagctcctc aggcacattta tgcggtaaat ggaatcgctg tccatcctgt
 1021 tcatggcacc cttgcaactg tgggatctga tggtagatc agcttctggg acaaagatgc
 1081 cagaacaaaa ctaaaaactt cggaacagtt agatcagccc atctcagctt gctgtttcaa
 1141 tcacaatgga aacatatttg catacgcttc cagctacgac tggatcaaagg gacatgaatt
 35 1201 ttataatccc cagaaaaaaa attacatttt cctgcgtaat gcagccgaag agctaaagcc
 1261 caggaataag aagtagtggc tggagactct ggctcagcca gagttgtttc tctccactct
 1321 gctcatctc tgtacgaatt tgggtcccag ccttggtggg ttgtcagcca tggacatgga
 1381 tttcaacccc tggagaaaac gatgtcattg ttcagcagct gagagcccag gcgtccgcgg
 1441 cgacttgccg tctctccatt ccactgcctg ttgcagagtt tttctgtaac taagggggtt
 40 1501 gaggttattg tagacgttag attgcgggca ccgccaggga ttttgcagcg cttcagtgta
 1561 cgtgttagag aatattggaa aagcgtctgt gagccccgtg ctgtattttg taataaagtc
 1621 ttttgcagat tgaaaaaaa aaaaaaaaaa

SEQ ID NO: 8

45 RNAExportHomol. (RAE1) U84720

/translation="MSLFGTTSFGFTSGTSMFGSATTDNHNPMKDIEVTSSPDDSIGC
 LSFSPPTLPGNFLIAGSWANDVRCWEVQDSGQTIPKAQQMHTGPVLDVCWSDDGSKVF
 TASC DKTA KMWD LSSNQAIQIAQHDAPVKTIHWIKAPNYSVMTGSDKTLKFW DTRS
 SNPMVVLQ LPERCYCADVIYPM AVVATAERGLIVYQLENQPS EFRRIESPLKHQHRCV
 50 AIFKDKNKPTGFALGSI EGRVAIH YINPPNPAKONFTFKCHRSNGTNTSAPQDIYAV
 NGIAFHPVHGTLATVGS DGRFSFWDKDARTKLTSEQLDQPI SACC FNHNGNIFAYAS
 SYDWSKGHEFYNPQKKNYIFLRNAAEELKPRNKK"

SEQ ID NO: 9

Bone morph. prot.7 (BMP7) BC008584

```

1 gggcgccagcg gggccccgtct gcagcaagtg accgaacggcc gggacggccg cctgccccct
5   61 ctgccacctg gggcggtgcg gggccggagc cggagcccg ggtagcgcgt agagccggcg
   121 cgatgcacgt gcgctcactg cgagctgcgg cgccgcacag cttcgtggcg ctctgggcac
   181 ccctgttcct gctgcgtccc gccctggccg acttcagcct ggacaacgag gtgcaactga
   241 gcttcatcca ccggcgccctc cgcagccagg agcggcgagg gatgcagcgc gagatcctct
   301 ccattttggg cttgccccac cgcccgccgc cgcacctcca gggcaagcac aactcggcac
  10   361 ccatgttcat gctggacctg tacaacgcca tggcggtgga ggagggcggc gggcccgccg
   421 gccagggcct ctcctacccc tacaaggccg tcttcagtac ccagggcccc cctctggcca
   481 gcctgcaaga tagccatttc ctcaccgacg ccgacatggt catgagcttc gtcaacctcg
   541 tggaacatga caaggaattc ttccacccac gctaccacca tcgagagttc cggtttgatc
   601 tttccaagat ccagaaggga gaagctgtca cggcagccga attccggatc tacaaggact
  15   661 acatccggga acgcttcgac aatgagacgt tccggatcag cgtttatcag gtgctccagg
   721 agcacttggg caggggaatcg gatctcttcc tgctcgacag ccgtaccctc tgggcctcgg
   781 aggagggctg gctgggtggtt gacatcacag ccaccagcaa ccactgggtg gtcaatccgc
   841 ggcacaacct gggcctgcag ctctcggtgg agacgctgga tgggcagagc atcaaccca
   901 agttggcggg cctgattggg cggcacgggc ccagaacaa gcagcccttc atggtggcct
  20   961 tcttcaaggc cacggaggtc cacttccgca gcatccggtc cacggggagc aaacagcgca
   1021 gccagaaccg ctccaagacg cccaagaacc aggaagccct gcggatggcc aacgtggcag
   1081 agaacagcag cagcgaccag aggcaggcct gtaagaagca cgagctgtat gtcagcttcc
   1141 gagacctggg ctggcaggac tggatcatcg cgcctgaagg ctacgccgcg tactactgtg
   1201 agggggagtg tgccttccct ctgaactcct acatgaacgc caccaaccac gccatcgtgc
   1261 agacgtggtt ccacttcata aacccggaac cggtgcccaa gccctgctgt gcgcccacgc
  25   1321 agctcaatgc catctccgtc ctctacttcg atgacagctc caacgtcctc ctgaagaaat
   1381 acagaaacat ggtgggtccg gcctgtggct gccactagct cctccgagaa ttcagaccct
   1441 ttggggccaa gtttttctgg atcctccatt gctcgccttg gccaggaacc agcagaccaa
   1501 ctgccttttg tgagaccttc cctccctat ccccaacttt aaaggtgtga gagtattagg
   1561 aaacatgagc agcatatggc ttttgatcag tttttcagtg gcagcatcca atgaacaaga
  30   1621 tcctacaagc tgtgcaggca aaacctagca ggaaaaaaaa acaacgcata aagaaaaatg
   1681 gccggggccg gtcattggct gggaagtcct agccatgcac ggactcgttt ccagaggtaa
   1741 ttatgagcgc ctaccagcca ggccaccag ccgtggggagg aagggggcgt ggcaaggggt
   1801 gggcacattg gtgtctgtgc gaaaggaaaa ttgaccggga agttcctgta ataaatgtca
      1861 caataaaacg aatgaatg

```

35

SEQ ID NO: 10

Bone morph. prot.7 (BMP7) BC008584

```

/translation="MHVRSLRRAAPHSFVALWAPLFLRLRSALADFSLDNEVHSSFIHR
RLRSQERREMQRILSLGLPHRPRPHLQGHNSAPMFMLDLYNAMAVEEGGGPGGQG
40  FSYPYKAVFSTQGPPLASLQDSHFLTDADMVMSFVNLVEHDKEFFHPRYHHREFRDL
SKIPEGEAVTAAEFRIYKDYIRERFDNETFRISVYQVLQEHLGRESDLFLLDSTLWA
SEEGWLVFDITATSNHWVNPRLNLGLQLSVETLDGQSIINPKLAGLIGRHGPQNKQPF
MVAFFKATEVHFIRSIRSTGSKQRSQNRSKTPKNQEALRMANVAENSSSDQRQACKKHE
LYVSFRDLGWQDWIIAPEGYAAYYCEGECAFLNSYMNATNHAIVQTLVHFINPETVP
45  KPCCAPTQLNAISVLVYFDDSSNVILKKYRNMVVRACGCH"

```

SEQ ID NO: 11

G-prot., α -stimul. (GNAS) BC002722

```

1 ccgcgcgcgc cgcagcccg cgcgcgcgcgc ccgcgcgcgc cgcgcgcgcgc cgcgcgcgcgc
50   61 ggaacagtaa gaccgaggac cagcgcaacg aggagaaggc gcagcgtgag gccaaacaaa
   121 agatcgagaa gcagctgcag aaggacaagc aggtctaccg ggccacgcac cgctgtctgc
   181 tgctgggtgc tggagaatct ggtaaaagca ccattgtgaa gcagatgagg atcctgcattg
   241 ttaatgggtt taatggagag ggcggcggaag aggacccgca ggctgcaagg agcaacagcg
   301 atgggtgagaa ggcaacccaa gtgcaggaca tcaaaaacaa cctgaaagag gcgattgaaa

```

5
10
15
20

```

361 ccattgtggc cgccatgagc aacctggtgc cccccgtgga gctggccaac cccgagaacc
421 agttcagagt ggactacatc ctgagtgtga tgaacgtgcc tgactttgac ttccctcccg
481 aattctatga gcatgccaaag gctctgtggg aggatgaagg agtgcgtgcc tgctacgaac
541 gctccaacga gtaccagctg attgactgtg cccagtactt cctggacaag atcgacgtga
601 tcaagcaggc tgactatgtg ccgagcgatc aggacctgct tcgctgccgt gtccctgactt
661 ctggaatctt tgagaccaag ttccagggtgg acaaagtcaa cttccacatg tttgacgtgg
721 gtggccagcg cgatgaacgc cgcaagtggg tccagtgcct caacgatgtg actgccatca
781 tcttcgtggg ggccagcagc agctacaaca tggatcatccg ggaggacaac cagaccaacc
841 gcctgcagga ggctctgaac ctcttcaaga gcatctggaa caacagatgg ctgcgaccca
901 tctctgtgat cctgttcctc aacaagcaag atctgctcgc tgagaaagtc cttgctggga
961 aatcgaagat tgaggactac tttccagaat ttgctcgcta cactactcct gaggatgcta
1021 ctcccagacc cggagaggac ccacgcgtga cccgggccaa gtacttcatt cgagatgagt
1081 ttctgaggat cagcactgcc agtggagatg ggcgctacta ctgctaccct catttcacct
1141 gcgctgtgga cactgagaac atccgccgtg tgttcaacga ctgccgtgac atcattcagc
1201 gcatgcacct tcgtcagtac gagctgctct aagaaggga cccccaatt taattaaagc
1261 cttaagcaca attaattaaa agtgaaacgt aattgtacaa gcagttaatc acccaccata
1321 gggcatgatt aacaaagcaa cctttccctt ccccgagtg attttgcgaa accccctttt
1381 ccttcagct tgcttagatg ttccaaattt agaaagctta aggcggccta cagaaaaagg
1441 aaaaaaggcc acaaaagttc cctctcactt tcagtaaaaa taaataaaac agcagcagca
1501 aacaaataaa atgaaataaa agaaacaaat gaaataaata ttgtgtgtg cagcattaaa
1561 aaaaatcaaa ataaaaatta aatgtgagca aag

```

SEQ ID NO: 12

G-prot., α -stimul. (GNAS) BC002722

25
30

```

/translation="MGCLGNSKTEDQRNEEKAQREANKKIEKQLQKDKQVYRATHRL
LLGAGESGKSTIVKQMRILHVNGFNGEGGEDPQAARSNSDGEKATKVQDIKNNLKEA
IETIVAAMSNLVPVELANPENQFRVDYILSVNVPDFDFPPEFYEHAKALWEDEGVR
ACYERSNEYQLIDCAQYFLDKIDVIKQADYVPSDQDLLRCRVLTSGIFETKFQVDKVN
FHMFDVGGQDERRKWIQC FNDVTAIIFVVAASSYNMVIREDNQTNRLQEALNLFKSI
WNNRWLRTISVILFLNKQDLLAEKVLAKGSKI EDYFPEFARYTT PEDATPEPGEDPRV
TRAKYFIRDEFRLISTASGDGRHYCYPHFTCAVD TENIRRVFNDCRDII QRMHLRQYE
LL"

```

SEQ ID NO: 13

35 Euk. Transl.init.2, subunit 2 (EIF2S2) BC000934

40
45
50
55

```

1 ggggtgtcgt ttcttttcgc tgatgcaaga gcctagtgcg gtgggtgggag aggtatcggc
61 agggggcagcg ctgccgcggg ggccctggggc tgaccgcgtc gacttcccgt ccgtgccgag
121 cccactcgag ccgcagccat gtctggggac gagatgattt ttgatccctac tatgagcaag
181 aagaaaaaga agaagaagaa gccttttatg ttagatgagg aaggggatac ccaaacagag
241 gaaaccagc cttcagaaac aaaagaagtg gagccagagc caactgagga caaggatttg
301 gaagctgatg aagaggacac taggaaaaaa gatgcttctg atgatctaga tgacttgaac
361 ttctttaatc aaaagaaaaa gaagaaaaaa actaaaaaga tatttgatat tgatgaagct
421 gaagaagggtg taaaggatct taagattgaa agtgatgttc aagaaccaac tgaaccagag
481 gatgaccttg acattatgct tggcaataaa aagaagaaaa agaagaatgt taagttccca
541 gatgaggatg aaatactaga gaaagatgaa gctctagaag atgaagacaa caaaaaagat
601 gatggtatct cattcagtaa tcagacaggc cctgcttggg caggctcaga aagagactac
661 acatacgagg agctgctgaa tcgagtgttc aacatcatga gggaaaagaa tccagatatg
721 gttgctgggg agaaaaggaa atttgtcatg aaacctccac aagtcgtccg agtaggaacc
781 aagaaaactt cttttgtcaa ctttacagat atctgtaaac tattacatcg tcagcccaaa
841 catctccttg catttttgtt ggtgaatttg ggtacaagtg gttctataga tggtataaac
901 caacttgtaa tcaaaggaa attccaacag aaacagatag aaaatgtctt gagaagatat
961 atcaaggaat atgtcacttg tcacacatgc cgatcaccgg acacaatcct gcagaaggac
1021 acacgactct atttcctaca gtgcgaaact tgtcattcta gatgttctgt tgccagtatc
1081 aaaaccggct tccaggctgt cacgggcaag cgagcacagc tccgtgccaa agctaactaa
1141 tttgctaate actgattttg caaagcttgt tgtggagatg tggctggaca ggtttgccat
1201 cagagtggat ataccgttgt attaaaaaca agataaaaaa gctgccaaaga tttttggcga

```

1261 gtggttggtc tgaagtcctt gcaagacgct gatgctcaag ctggtgacat actcattgcc
1321 tactttaaca cctgtcagag aaacgtgata tggggtaagg aggtgctttt ttaaaatcgt
1381 tcatagactt ctgtaaaatg caagataaat taaagttatt ataacagtga ttctttcaa

5 SEQ ID NO: 14

Euk. Transl.init.2, subunit 2 (EIF2S2) BC000934

/translation="MSGDEMI F D P T M S K K K K K K K K P F M L D E E G D T Q T E E T Q P S E T K E V
E F G P T E D K D L E A D E E D T R K K D A S D D L D L N F F N Q K K K K K K T K K I F D I D E A E G V K D L K
I E S D V Q E P T E P E D D L D I M L G N K K K K K K N V K F P D E D E I L E K D E A L E D E D N K K D D G I S F S
10 M Q T G P A W A G S E R D Y T Y E E L L N R V F N I M R E K N P D M V A G E K R K F V M K P P Q V V R V G T K K T S
F V N F T D I C K L L H R Q P K H L L A F L L A E L G T S G S I D G N N Q L V I K G R F Q Q K Q I E N V L R R Y I K
E Y V T C H T C R S P D T I L Q K D T R L Y F L Q C E T C H S R C S V A S I K T G F Q A V T G K R A Q L R A K A N"

SEQ ID NO: 15

15 Dynein lt. Chain A2 (DNCL2A) NM_014183

1 cgcagaaagg cacaggactc gctaagtgtt cgctacgcgg ggctaccgga tcggctcgaa
61 atggcagagg tggaggagac actgaagcga ctgcagagcc agaagggagt gcaggggaatc
121 atcgtcgtga acacagaagg cattcccatc aagagcacca tggacaaccc caccaccacc
181 cagtatgccg gcctcatgca cagcttcacg ctgaaggcac ggagcaccgt gcgtgacatc
20 241 gacccccaga acgatctcac cttccttcga attcgctcca agaaaaatga aattatgggt
301 gcaccagata aagactatct cctgattgtg attcagaatc caaccgaata agccactctc
361 ttggctccct gtgtcattcc ttaatttaat gcccccaag aatgttaatg tcaatcatgt
421 cagtggacta gcacatggca gtcgcttgga acccactcac accaatccag tgaccgtgtg
481 tgggctggcg gctcttctcc cccaccaacg gaacccctgt gtgcaccaac cttccccaga
25 541 gctccggagc gccctctcct cacttccagg ttttgagca agagcttgca ggaagccgc
601 acccagcttc cttctgacct tcagttcact ttgtcgccct tggagaaagc tgtttttctt
661 taactaaaaa taaccaaaat gcttaaaaaa aaaaaaaaaa aa

SEQ ID NO: 16

30 Dynein lt. Chain A2 (DNCL2A) NM_014183

/translation="MAEVEETLKR LQSQKGVQGIIVNTEGIPIKSTMDNPTTTQYAS
LMHSFILKARSTVRDIDPQNDLTFLRIRSKKNEIMVAPDKDYFLIVIQNPTE"

SEQ ID NO: 17

35 Proteosome subunit α -7 (PSMA7) BC004427

1 cggcgccgag ggtggggcgc gggcgtagtg gcgcggggag tcgcgggtgc gcgcggggccg
61 tgagtgtgcg cttttgagag tcgcggcgga aggagcccgg ccgcccggcg ccggcatgag
121 ctacgaccgc gccatcaccg tcttctcgcc cgacggccac ctcttccaag tggagtacgc
181 gcaggaggcc gtcaagaagg gctcgaccgc ggttgggtgt cgaggaagag acattgttgt
40 241 tcttgggtgtg gagaagaagt cagtggccaa actgcaggat gaaagaacag tgcggaagat
301 ctgtgctttg gatgacaacg tctgcatggc ctttgcaggc ctcaccgccc atgcaaggat
361 agtcatcaac agggcccggg tggagtgcca gagccaccgg ctgactgtgg aggaccgggt
421 cactgtggag tacatcaccg gctacatcgc cagtctgaag cagcgttata cgcagagcaa
481 tgggcgccagg ccgtttggca tctctgccct catcgtgggt ttcgactttg atggcactcc
45 541 taggctctat cagactgacc cctcgggcac ataccatgcc tgggaaggcca atgccatagg
601 tcgggggtgcc aagtcatgac gcgagttcct ggagaagaac tatactgacg aagccattga
661 aacagatgat ctgaccatta agctggtgat caaggcactc ctggaagtgg ttcagtcagg
721 tggcaaaaac attgaacttg ctgtcatgag gcgagatcaa tccctcaaga ttttaaatcc
781 tgaagaaatt gagaagtatg ttgctgaaat tgaaaaagaa aaagaagaaa acgaaaagaa
50 841 gaaacaaaag aaagcatcat gatgaataaa atgtctttgc ttgtaatttt taaattcata
901 tcaatcatgg atgagtctcg atgtgtaggc ctttccattc catttattca cactgagtgt

961 cctacaataa acttccgtat tttt

SEQ ID NO: 18

Proteosome subunit α -7 (PSMA7) BC004427

5 /translation="MSYDRAITVFSPDGHLFQVEYAQEA VKKGSTAVGVRGRDIVVLG
VEKKSVAKLQDERTVRKICALDDNVCMAFAGLTADARIVINRARVECSHRLTVEDPV
TVEYITRYIASLKQRYTQSNRRPFGISALIVGFDFDGTPLRYQTDPSGTYHAWKANA
IGRGAKSVREFLEKNYTDIAETDDLTIKLVIKALLEVVQSGGKNIELAVMRDQSLK
ILNPEEIEKYVAEIEKEKEENEKKKQKKAS"

10

SEQ ID NO: 19

Activity dep. Neuroprotector (ADNP) AF250860

1 cggccgcggc gcgagccgga gtccgcccag ccggagcgcg acgaggcccc gggcgcgccc

15 61 tccccgctgc cgccaccgcc gtgcgcgcgc catccgcccc cgccgcgcgc cgctgtccgg
121 cccccgagca cgccggcccc gcgcgcgcct cgaggccgag tcaaggtgtg agatgcacaa
181 tgcgaaacct aggccccagc ttttacacca tgatgcgcag ggttgactt tttgtactga
241 actgataggt ggcctagtgg ttatgccctg tactaccatt ttgaggatct ggactccgtt
301 tctgccttg ctctttggac cacattgtca attcacaccg aaactatgtt ccaacttctt
361 gtcaacaatc ttggcagttt aagaaaagcc cggaaaactg tgaaaaaat acttagtgac

20 421 attgggttg aatactgtaa agaacaata gaagatttta acaatttga acctaatac
481 ttttatttga aaaacactac atgggaggat gtaggactgt gggaccatc acttacgaaa
541 aaccaggact atcgacaaa acctttctgc tgcagcgtt gtccatttct ctaaaaattc
601 ttctctgcct acaaaagtca tttccgcaat gtccatagt aagactttga aaataggatt
661 ctctttaatt gcccctactg taccttcaat gcagacaaaa agacttttga aacacacatt

25 721 aaaatatttc atgctccgaa cgccagcgca ccaagtagca gcctcagcac tttcaaagat
781 aaaaacaaaa atgatggcct taaacctaa caggctgaca gtgtagagca agctgtttat
841 tactgtaaga agtgcaacta ccgagatcct ctttatgaaa tagttaggaa gcacatttac
901 agggaaacatt ttcagcatgt ggcagcacct tacatagcaa aggcaggaga aaaatcactc
961 aatggggcag tccccttagg ctccaatgcc cgagaagaga gtagtattca ctgcaagcga

30 1021 tgccttttca tgccaaagtc ctatgaagct ttggtacagc atgtcatcga agaccatgaa
1081 cgtataggct atcaggtcac tgccatgatt gggcacacaa atgtagtggg tccccgatcc
1141 aaacccttga tgctaattgc tcccaaacct caagacaaga agagcatggg actcccacca
1201 aggatcgggt cccttgcttc tggaaatgtc cggcttttac catcacagca gatggtgaat
1261 cgactctcaa taccaaagcc taacttaa atctacaggag tcaacatgat gtccagtgtt

35 1321 catctgcagc agaacaacta tggagtcaaa tctgtaggcc agggttacag tgttggtcag
1381 tcaatgagac tgggtctagg tggcaacgca ccagtttcca ttctcaaca atctcagtct
1441 gtaaagcagt tacttccaag tggaaacgga aggtcttatg ggcttgggtc agagcagagg
1501 tcccaggcac cagcaagata ctccctgcag tctgcta atg cctctctct ctcactcggc
1561 cagttaaagt ctccctccct ctctcagtc caggcatcca gactgttagg tcagtccagt

40 1621 tccaaacctg ctgcagctgc cacaggccct ccccaggta acacttctc aactcaaaag
1681 tggaaaatat gtacaatctg taatgagctt tttctgaaa atgtctatag tgtgcacttc
1741 gaaaaagaac ataaagctga gaaagtcca gcagtagcca actacattat gaaaatacac
1801 aattttacta gcaa atgcct ctactgta at cgctatttac ccacagatac tctgctcaac
1861 catatgttaa tcatggtct gtctgtcca tattgcccgt caactttcaa tgatgtggaa

45 1921 aagatggccg cacacatgag gatggttcac attgatgaag agatgggacc taaaacagat
1981 tctactttga gttttgattt gacattgcag cagggtagtc aactaactat ccatctcctg
2041 gtaactacat acaatctgag ggatgcccc gctgaatctg ttgcttacca tgccccaaat
2101 aatcctccag ttctccaaa gccacagcca aagggttcagg aaaaaggcaga tatccctgta
2161 aaaagttcac ctcaagctgc agtgccctat aaaaaagatg ttgggaaaac cctttgtcct

50 2221 ctttgccttt caatcctaaa aggaccata tctgatgcac ttgcacatca cttacgagag
2281 aggaccaag ttattcagac ggttcattca gttgagaaaa agctcaccta caaatgtatc
2341 cattgccttg gtgtgtatac cagcaacatg accgcctcaa ctatcactct gcactagtt
2401 cactgcaggg gcgttgga aa gacccaaaat ggccaggata agacaaatgc accctctcgg
2461 cttaatacagt ctccaagtct ggcacctgtg aagcgcactt acgagcaaat ggaatttccc

55 2521 ttactgaaaa aacgaaagt agatgatgat agtgattcac ccagcttctt tgaagagaag
2581 cctgaagagc ctgttgtttt agcttttagac cccaagggtc atgaagatga ttcctatgaa

2641 gccaggaaaa gcttttctaac aaagtatttc aacaaacagc cctatcccac caggagagaa
 2701 attgagaagc tagcagccag tttatgggta tggaagagtg acatcgcttc ccatttttagt
 2761 aacaaaagga agaagtgtgt ccgtgattgt gaaaagtaca agcctggcgt gttgctgggg
 5 2821 ttttaacatga aagaattaaa taaagtcaag catgagatgg attttgatgc tgagtgggcta
 2881 tttgaaaatc atgatgagaa ggattccaga gtcaatgcta gtaagactgc tgacaaaaag
 2941 ctcaaccttg ggaaggaaga tgacagttcc tcagacagtt ttgaaaattt ggaagaagaa
 3001 tccaatgaaa gtggtagccc ttttgaccct gtttttgaag ttgaacctaa aatctctaac
 3061 gataaccagc aggaacatgt actgaaggta attcctgagg atgcttcaga atctgaggag
 10 3121 aagctagacc aaaaagagga tggttcaaaa tacgaaacta ttcatttgac tgaggaacca
 3181 accaaactaa tgcacaatgc atctgatagt gaggttgacc aagacgatgt tgttgagtgg
 3241 aaagacgggtg cttctccatc tgagagtggg cctggatccc aacaagtgtc agactttgag
 3301 gacaatacct gcgaaatgaa accaggaacc tggctctgac agtcttccca aagcgaagat
 3361 gcaaggagca gtaagccagc tgccaaaaaa aaggctacca tgcaagggtga cagagagcag
 3421 ttgaaatgga agaatagttc ctatggaaaa gttgaagggt tttgggtctaa ggaccagtca
 15 3481 cagtgggaaga atgcatctga gaatgatgag cgcttatcta acccccagat tgagtggcag
 3541 aatagcaciaa ttgacagtga ggatggggaa cagtttgaca acatgactga tggagtagct
 3601 gagcccatgc atggcagctt agccggagtt aaactgagca gccaacaggc ctaagtgccca
 3661 ggttccctgg cgttggtgac atgctgcagc ctggaactct gatctccagt gtgactgcaa
 20 3721 agctgtcttc tcaactggtac tgccttgtga gtactgggtg gactgtgggg catgtggcgg
 3781 ctgcagttcc agtggttatt tctaagtcta tgacaggaca ggctgttctt gcttcagaac
 3841 cttctctgac agacacggtg actaaatgtg aaaaaccaat aagctgggtga ctcatgaata
 3901 cacacgagga aaagcagagg tttattttat ctgccttttc aacatttctt tccctctgtg
 3961 aaatgattgg tcagatgtct ttgagaagtg ttaaactaat tcacatggta gtgtagggcc
 25 4021 aacatacaag ctaccagtct aatgtgtata gtagactttg ggaaaagcga ttttttttca
 4081 tgtattcatt ctgaatagtt gaaatgtata tttgtacagt ctttttagacc tattcaagtg
 4141 atgctcatga tcctgttact gtgtgcccac catagatttc ttttttttagt gttgcccttg
 4201 ctgtgtaata aacgctctat ctagtttacc tagcaaaagc tcaaaactgc gctagtagtg
 4261 acttttttga cagacttagt ttttgacat aaccttgtag aatcttgcaa cagagccag
 30 4321 ccacgtaaga tatatatctg gactctcttg tattatagga tttttcttgt tctgaatata
 4381 cttgacatta cagctgtcaa aaacaaaaaac tggattttca gatctgtttt ctgaaatctt
 4441 ttaagctaaa atcacatgca agaattgact ttgcagctac taattttgac accttttaga
 4501 tctgtataaa agtgtgttgt gttgaagcag caaaccaatg agtgctgcat tttggatatt
 4561 tagttttatc ttttagttcaa caccatcatg gtggattcat ttataccatc taatatatga
 4621 cacactgttg tagtatgtat aatttttgtga tctttatttt ccctttgtat tcattttaag
 35 4681 catctaaata aattgctgta ttgtgcttaa tgt

SEQ ID NO: 20

Activity dep. Neuroprotector (ADNP)

AF250860

/translation="MFQLPVNNLGLSLRKARKTVKKILSDIGLEYCKEHIEDFKQFEPN
 40 DFYLNKNTTWEDVGLWDPSTLTKNQDYRTKPFCCSACPFSSKFFSAYKSHFRNVHSEDFE
 NRILLNCPYCTFNADKKTLETHIKIFHAPNASAPSSSLSTFKDKNKNDGLKPKQADSV
 EQAVYYCKKCTYRDPPLYEIVRKHIYREHFQHVAAPIYAKAGEKSLNGAVPLGSNAREE
 SSIHCKRCLFMPKSYEALVQHVIEDHERIGYQVTAMIGHTNVVVRSPKPLMLIAPKPQ
 45 DKKSMGLPPRIGSLASGNVRSLSQQMVRNLSIPKPNLNSTGVNMMSSVHLQQNNYGV
 KSVGQGYSVGQSMRLGLGGNAPVSIQQSQSVKQLLPSGNRSYGLGSEQRSQAPARY
 SLQSANASSLSGQLKSPSLSQSASRVLGQSSSKPAAATGPPPGNTSSTQKWKICT
 ICNELFPENVYSVHFEKEHKAKEVPAVANYIMKIHNFTSKCLYCNRYLPTDTLLNHML
 IHGLSCP YCRSTFNDVEKMAAHMRMVHIDEEMGPKTDSTLSFDLTLQQGSHTNIHLLV
 50 TTYNLRDAPAESVA YHAQNPPVPPKPQPKVQEKADIPVKSSPQAAVPYKKDVGKTLCL
 PLCFSILKGPISDALAHHLRERHQVIQTVHPVEKKLTYKCIHCLGVYTSNMTASTITL
 HLVHCRGVGKTQNGQDKNAPSRLNQSPLAPVKRTYEQMEFPLLLKKRKLDDSDSPS
 FFEEKPEEPVVLALDPKGHEDDSYEARKSFLT KYFNKQPYPTREIEKLAASLWLWKS
 DIASHFSNKRKKCVRDCEKYKPGVLLGFNMKELNKVKHEMDFDAEWLFENHDEKDSRV
 55 NASKTADKKLNLGKEDDSSDSFENLEESNESGSPFDPVFEVEPKISNDNPPEHVLK
 VIPEDASESEEKLDQKEDGSKYETIHLTEEPTKLMHNASDSEVDQDDVVEWKDGASPS
 ESGPGSQQVSDFEDNTCEMKPGTWSDESSQSEDARSSKPAAKKKATMQGDREQLKWKV
 SSGKVEGFWKQSQWKNASENDERLSNPQIEWQNSTIDSEGEQFDNMTDGVAEPM
 HGLAGVKLSSQQA"

SEQ ID NO: 21

C20orf129 AK055793

```
5      1 aaaaagcagc caatgggaga gccgaggcgg ggaggtgcgg ccaatggcgc gggcctgttt
      61 gattcaaagg ttgcctataa agcgggactg cacgccgggt tttgtccgag ggctgtcgag
     121 tccgagcgcc gccatggctc tgctgtccga gggcctggac gaggtgcccg ccgctgcct
     181 gtcgccgtgc gggccgcccc acccgaccga gctgttcagc gagtcacggc gcctggctct
     241 ggaggagctg gtggcgggcg gccccgaagc cttcgcgggc ttcctgcgac gcgagcgctt
10    301 ggctcgtttc ctgaaccccg atgagggtga cgccattctg cgcgcggcgg agaggccggg
     361 agaggagggc gcggcgggcg cgccggcgcg cgaggactcg ttcggctcct cgcacgactg
     421 ctcttcgggc acctacttcc ccgagcagtc ggacctggag ccaccgctgt tggagcttgg
     481 ctggcccgcc ttctaccagg gcgcctaccg cggcgccacg cgtgtcgaga cgcacttcca
     541 gccccgcggc gctggcgaag gtggccccta cggctgcaag gacgctctgc gccagcagct
15    601 ccgctcggcg cgagaggtga ttgcagtggg catggacgtg ttcacagaca tcgacatctt
     661 cagagacctg caagaaatat gcaggaaaaca gggagtgtgt gtgtatatcc ttctggacca
     721 ggctctcctc tctcaatttc tggatatgtg catggatctg aaagttcata ctgaacagga
     781 aaagttaatg acagttcgga ctatcacagg aaatatctac tatgcaaggt caggaactaa
     841 gattattggg aaggttcacg aaaagttcac gttgattgat ggcacccgag tggcaacagg
20    901 ctctacagtc tttacataga cggatggcaa attaaacagc agtaacttgg taattctgtc
     961 tggccaagtg gttgaacact ttgatctgga gttccgaatc ctgtatgccc agtccaagcc
    1021 catcagcccc aaactcctgt ctcaacttcca gagcagcaac aagtttgatc acctcaccaa
    1081 ccgaaaacca cagtccaagg agctcaccct gggcaacctg ctgcggatgc ggctggctag
    1141 gctgtcaagt actcccagga aggcggacct ggaccagag atgcccgcag agggcaaggc
25    1201 agagcgcaag ccccatgact gtgagtcctc tactgttagt gaggaagact acttcagcag
    1261 ccacagggac gagctccaga gcagaaaggc cattgacgct gccactcaaa cagagccagg
    1321 agaggagatg ccagggtgta gtgtgagtga ggtgggaaca caaaccagca tcaccacagc
    1381 atgtgctggg acccagactg cagtcatcac caggatagca agctctcaaa ccacgatttg
    1441 gtccagatcg accactactc agactgacat ggatgagaac attctcttcc ctcgaggaac
30    1501 tcaatctaca gaagggtcac cagtctcaaa aatgtctgta tcgagatctt ccagtttgaa
    1561 gtcttctctc tctgtgtctt cccaaggctc tgtggcaagc tccactgggt ctcccgttcc
    1621 catcagaacc actgacttcc acaatcctgg ctatcccaag tacctgggca cccccacct
    1681 ggaactgtac ttgagtgact cacttagaaa cttgaacaaa gagcggcaat tccacttcgc
    1741 tggatatcagg tcccggctca accacatgct ggctatgctg tcaaggagaa cactctttac
35    1801 tgaaaaccac cttggccttc attctggcaa tttcagcaga gttaatttgc ttgctgttag
    1861 agatgtagca ctttatcctt cctatcagta actgctccgt gttcagactc ctggtttctt
    1921 ccaggcttac agtggacatc atcagcttcc tgctttaaaa aatatcttat gctgcataat
    1981 gcctttcttt taactgactt tgtcaccttt gttgtctttg aattctttag gctgcataat
    2041 attttacatg ctttgttttg tcatgtatat accaggtatt ggttttatgg tttaaacact
40    2101 atggatacag gggtttgttt tgcacaattt taatagtcac gcactacata atgatgtttt
    2161 ggtcaatgac agaccacgta tatgttggca gtctcataag attataatac tgtattttta
    2221 ctataccttt tctgtgttta gatacaaata ccattatgtt acagttgcct acagtattca
    2281 gtgcagtaac atgatgtaca ggtttgtagc ctgttttgca tttttcttag gttgtatgct
    2341 cttctgtttt aaaggtttga atcaccagca tttttgtgat caaaatccta tttagaaaaa
45    2401 ataaaactac tttctgttta tctcttt
```

SEQ ID NO: 22

C20orf129 AK055793

```
50 /translation="MARACLIQRLPIKRDCTPVFVRGLSSPSAAMALLSEGLDEVPAALSPCGPPNPTELFSESRRLLALEELVAGGPEAFAAFLRRERLARFLNPDEVHAILRAA
ERPGEEGAAAAAAEDSFGSSHDCSSGTYFPEQSDLEPPLLELWPAFYQGAYRGATR
VETHFQPRGAGEGGPYGCKDALRQQLRSAREVIAVMDVFTDIDIFRDLQEICRKQGV
AVYILLDQALLSQFLDMCMDLKVHPEQEKLMTVRTITGNIYYARSGTKIIGKVHEKFT
LIDGIRVATGSYSFT"
55
```

SEQ ID NO: 23

C20orf52 BC008488

```
1 gacgcggggc cggaacgcga agaggggtgt ggagtcgggc taccactga ttttcttcc
5 61 cttacttccc ctgagccctt gggcccactt cccagcctac cgcttccgtc cccgcccagc
121 tcttgggcca gcgcctgggc ccacactttc ctatcccccg cagatgccgg tggccgtggg
181 tccctacgga cagtccagc caagctgctt cgaccgtgtc aaaatgggct tcgtgatggg
241 ttgcgcctgt ggcattggcg cggggcgctt ctccggcacc ttttctgtc tcaggatcgg
301 aatgcggggt cgagagctga tgggcggcat tgggaaaacc atgatgcaga gtggcggcac
10 361 ctttggcaca ttcattggcca ttgggatggg catccgatgc taacctggt tgccaactac
421 atctgtccct tcccatcaat cccagcccat gtactaataa aagaaagtct ttgagtaaaa
481 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa
541 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa
601 aa
```

15 SEQ ID NO: 24

C20orf52 BC008488

/translation="MPVAVGPYQSQPSCFDRVKMGFVMGCAVGMAAGALFGTFSLR
IGMRGRELMMGGIGKTMQSGGTFGTFMAIGMGIRC"

20 SEQ ID NO: 25

C20orf20 BC018841

```
1 ctcccgccgg gggctccttg ctggccggg ccgcccggcat gggagaggcc gaggtgggag
61 gcggggggcg cgcaggcgac aagggcccg gggaggcgcc caccagccc ggcggaggaga
25 121 cagtgggtgt gagccccgag gtggagggtg gcctcttcca cgccatgctg ggccacaagc
181 ccgtcgggtg gaaccgacac ttccacatga ttgtattcg ggacaagttc agccagaaca
241 tcgggcggca ggtcccatcc aaggtcatct gggaccatct gagcaccatg tacgacatgc
301 aggcgctgca tgagtctgag attcttccat tcccgaatcc agagaggaaac ttcgtccttc
361 cagaagagat cattcaggag gtccgagaag gaaaagtgat gatagaagag gagatgaaag
421 aggagatgaa ggaagacgtg gacccccaca atggggctga cgatgttttt tcatcttcag
30 481 ggagtttggg gaaagcatca gaaaaatcca gcaaagacaa agagaagaac tctcagact
541 tggggtgcaa agaaggcgca gacaagcgga agcgagccg ggtcaccgac aaagtcctga
601 ccgcaaacag caacccttcc agtcccagtg ctgccaaagc gcgccgcacg tagaccctca
661 gccctggttg cggcagagaa gcgggcgagg cactgtggtc gctgaggggg ttggttgggt
721 ctgagtggca cccccaggc acagtgtatc catcccagtg ccatgagccc acactgcccg
35 781 ccctcaggct ctcaggtgaa cgtggccgtc agcggggaaa cgtgtgtgtc agttggacca
841 tgtgggaccc tgatggacct gaaagaccag gatcgggtcca gctcagatat tgagggctct
901 gaagcctagt tctgtcttct ctggagcagc tgtggcttcc ccgtggctgc ttggtgacat
961 ggattagcgc tacgtgggct gcagcatttg ggatccaggc tacctagagg ggcattgggc
40 1021 cagggaaaac ctccgattag caagcaataa aaacatgacc tcaactctcc tcaaaggagc
1081 ccctggtctt ccctgtgtga ctcagttctt tccatctgtt tgtcccgtg caagcctctt
1141 tctgcgctga ctgtgacatt ggaacgtggc ctctctgtca cccctccgt gccacgcact
1201 gaaggccacc cccaccacc tgggaaacta agaactggat attttgctc attcacttgt
1261 actgtaacaa tgtatataat ttggttggtt ttctactatt taatttttaa gaagcctatt
45 1321 ttactagtgt tttatatgaa caaagtactg cagaagttaa acctgtgttg tattttttct
1381 gagatgtttt gctttaagag atactttttg ctgagttttt atatgccaga tacagagaat
1441 ttgtagcggg tatttttgta tgatctagta acttgcaaac agaccaaag gatgagaggc
1501 ggggaccgtg cagctgtcgg ctgatgagga ggcggccgcc ccagtgtgta tggagatgcc
1561 actttcgtgt gactgcgaac attaaagcac aaaaaaaatc caaaaaaaaa aaaaaaaaaa
50 1621 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa
1681 aaaaa
```

SEQ ID NO: 26

C20orf20 BC018841

/translation="MGEAEVGGGGAAGDKGPGEAATSPAETTVVWSPEVEVCLFHAML
GHKPVGVNRHFHMICIRDKFSQNIGRQVPSKVIWDHLSTMYDMQALHESEILPFPNPE
RNFVLPEEIIQEVREGKVMIEEEMKEEMKEDVDPHNGADDVFSSSGSLGKASEKSSKD
KEKNSSDLGCKEGADKRKRSRVTDKVL TANSNPSSPSAAKRRRT"

SEQ ID NO: 27

C20orf188 BC013144

1 gacatggcgg cggcgccgggt agcggctggg tctggagccg gccgagggag acggtcggca
61 gccacagtgg cggcttgggg cggatggggc ggccggccgc ggcctggtaa cattctgctg
121 cagctgcggc agggccagct gaccggccgg ggcctgggcc gggcggtgca gttcactgag
181 acttttttga cggagaggga caaacatcc aagtggagtg gaattcctca gctgctcctc
241 aagctgcaca ccaccagcca cctccacagt gactttgttg agtgtaaaaa catcctcaag
301 gaaatttctc ctcttctctc catggaggct atggcatttg ttactgaaga gaggaaactt
361 acccaagaaa ccacttatcc aaatacttat atttttgact tgtttgaggg tgttgatctt
421 cttgtagaaa ttcttatgag gcctacgata tctatccggg gacagaaact gaaaataagt
481 gatgaaatgt ccaaggactg cttgagtatc ctgtataata cctgtgtctg tacagagggg
541 gttacaaagc gtttggcaga aaagaatgac tttgtgatct tcctgtttac attgatgaca
601 agtaagaaga cattcttaca aacagcaacc ctcattgaag atattttggg tgtaaaaaag
661 gaaatgatcc gactagatga agtccccaat ctgagttcct tagtatccaa ttctgatcag
721 cagcagctcg ctaatttctg cgggattctg gctgtcacca tttcagagat ggatacaggg
781 aatgatgaca agcacacgct tcttgccaaa aatgctcaac agaagaagag cttgagtttg
841 gggccttctg cagctgaaat caatcaagcg gcccttctca gcattcctgg ctttgttgag
901 cggctttgca aactggcgac tcgaaagggt tcagagtcaa cgggcacagc cagcttcctt
961 caggagttgg aagagtggta cacatggcta gacaatgctt tgggtgctaga tgccctgatg
1021 cgagtggcca atgaggagtc agagcacaa caagcctcca ttgtgttccc tcctccaggg
1081 gcttctgagg agaattggcct gcctcacacg tcagccagaa cccagctgcc ccagtcaatg
1141 aagattatgc atgagatcat gtacaaactg gaagtgtct atgtcctctg cgtgctgctg
1201 atggggcgct agcgaaacca ggttcacaga atgattgcag agttcaagct gatccctgga
1261 cttaataatt tgtttgacaa actgatttgg aggaagcatt cagcatctgc ccttgtcctc
1321 catggtcaca accagaactg tgactgtagc cgggacatca ccttgaagat acagtttttg
1381 aggtctcttc agagcttcag tgaccaccac gagaacaagt acttggtact caacaaccag
1441 gagctgaatg aactcagtgc catctctctc aaggccaaca tccttgaggt ggaagctgtc
1501 ctcaacaccg acaggagtgt ggtgtgtgat gggaagaggg gcttattaac tcgtctgctg
1561 caggctcatg agaaggagcc agcagagtcg tctttcaggt tttggcaagc tcgggctgtg
1621 gagagtttcc tccgagggac cacctcctat gcagaccaga tgttcctgct gaagcgaggg
1681 ctcttgagag acatccttta ctgcattgtg gacagcgagt gtaagtcaag ggatgtgctc
1741 cagagttact ttgacctcct gggggagctg atgaagttca acgttgatgc attcaagaga
1801 ttcaataaat atatcaacac cgatgcaaag ttccaggtat tcctgaagca gatcaacagc
1861 tccctgggtg actccaacat gctggtgcgc tgtgtcactc tgtccctgga ccgatttgaa
1921 aaccaggtgg atatgaaagt tgccgaggta ctgtctgaat gccgctgct cgccacata
1981 tccaggtgac ccacgcagat gtccttctc ttcgcctca tcaacatcat ccacgtgcag
2041 acgctgaccc aggagaacgt cagctgcctc aacaccagcc tggatgacct gatgctggcc
2101 cgacggaaag agcggctgcc cctgtacctg cggctgctgc agcggatgga gcacagcaag
2161 aagtaccccg gcttcctgct caacaacttc cacaacctgc tgcgcttctg gcagcagcac
2221 tacctgcaca aggacaagga cagcacctgc ctagagaaca gtcctgcat cagcttctca
2281 tactggaagg agacagtgtc catcctgttg aaccgggacc ggcagtcacc ctctgctctc
2341 gttagctaca ttgaggagcc ctacatggac atagacaggg acttactga ggagtgcact
2401 tgggcccagg ctcgggaggg tgctgggcca gtgtgggtga gcgtgggtac gatgccacac
2461 gccctgcctt gttcccgctt cctctgctg cctctgctc gccccaggtc tttgggtaca
2521 ggcttggtgg gagggagtc ctagaagccc ttgggtcccc tgggtctgag ggccctaggt
2581 catggagagc ctcagtcctc ataattgagga cagggtacca tgcccacctt tccttcagaa
2641 cctgggggcc cagggccacc cagaggtgag aggacattta gcattagctc tgtgtgagct
2701 cctgccggtt tcttggtgtg cagtcagtc cagagtgggg aggaagatat ggggtgaccc
2761 ccccccccat ctgtgagcca agcctccctt gtccctggcc tttggaccca ggcaagggt
2821 tctgagccct gggcaggggt ggtgggtacc agagaatgct gccttcccc aagcctgccc

2881 ctctgcctca ttttcctgta gctcctctgg ttctgtttgc tcattggctg ctgtgttcat
2941 ccaagggggg tctcccagaa gtgagggggc ttccctcca tcccttgagg cacggggcag
3001 ctgtgcctgc cctgcctctg cctgaggcag ccgctcctgc ctgagcctgg acatggggcc
3061 cttccttggtg ttgccaattt attaacagca aataaaccaa ttaaatggag actattaaat
3121 aactttattt taataaaaaa aaaaaaaa

SEQ ID NO: 28

C20orf188

BC013144

/translation="MAAAPVAAĠSGAGRGRRSAATVAAWGGWGGRRPRPGNILLQLRQG
10 QLTGRGLVRAVQFTETFLTERDKQSKWSGIPQLLLKLHTTSHLHSDFVECONILKEIS
PLLSMEAMAFVTEERKLQETTYPNTYIFDLFGGVDLLVEILMRPTISIRGQKLKISD
EMSKDCLSILYNTCVCTEGVTKRLAEKNDFVIFLFTLMTSKKTFLQTATLIEDILGVK
KEMIRLDEVPNLSSLVSNFDQQQLANFCRILAVTISEMDTGNDKHTLLAKNAQQKKS
LSLGPSAAEINQAALLSIPGFVERLCKLATRKVSESTGTASFLQELEEWYTWLDNALV
15 LDALMRVANESEHNQASIVFPPPGASEENGLPHTSARTQLPQSMKIMHEIMYKLEVL
YVLCVLLMGRQRNQVHRMIAEFKLIPLNNLFDKLIWRKHSASALVLHGHNNQNCDCSP
DITLKIQFLRLQLQSFSDDHKNYLLNNQELNELSAISLKANIPVEEAVLNTDRSLVC
DGKRGLLTRLLQVMKKEPAESSFRFWQARAVESFLRGTTSYADQMFLKRGLEHILY
CIVDSECKSRDVLQSYFDLLGELMKFNVDFAKRFNKYINTDAKFQVFLKQINSSLVDS
20 NMLVRCVTLSLDRFENQVDMKVAEVLSECRLLAYISQVPTQMSFLFRLINIIHVQTLT
QENVSCLNNTSLVILMLARRKERLPLYLRLLQRMESKKYPGFLNNFHNLLRFWQQHY
LHKDKDSTCLENSSCISFSYWKETVSILLNPDRQSPSALVSYIEEPYMDIDRDFTEE"

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.